

Exercise improves anxiety and mood in older adults undergoing chemotherapy

July 3 2019

Although we know that exercise improves anxiety and mood problems in younger people with cancer, few studies have looked at the effects of exercise on older adults with cancer. Since most new cancer cases occur in adults aged 60 or older, a team of researchers from the University of Rochester Medical Center and other institutions designed a study to learn more.

Their study appeared in the June issue of the *Journal of the American Geriatrics Society (JAGS)*.

Having cancer increases the chances of people experiencing anxiety and mood issues, which can affect emotional and social well-being. In turn, this may lead people to discontinue cancer treatments—which can mean shortening their survival.

Chemotherapy can benefit older adults with cancer, even though [older people](#) receiving this type of treatment often experience higher rates of dangerous side effects than [younger people](#) do. Older adults often experience anxiety and other [mood disorders](#) during their treatment for cancer, too—and treating those problems with medications can often cause potentially dangerous side effects.

What's more, many anti-anxiety medications such as benzodiazepines and antidepressants are listed in the American Geriatrics Society (AGS) Beers Criteria as being potentially inappropriate for older adults. That's why it is desirable to seek alternative treatments that are safe and

effective at improving anxiety, mood disturbances, and emotional and social well-being, including treatments that don't rely on medications. For example, several studies have been conducted to examine the relationship between exercise and mood in cancer survivors and most have shown positive results.

The researchers in the new *JAGS* study examined the Exercise for Cancer Patients (EXCAP) program, a home-based, low- to moderate-intensity aerobic and resistance exercise program. In the study, those who were assigned to the EXCAP program received an exercise kit. It contained a pedometer, three exercise bands (medium, heavy, extra heavy), and an [instruction manual](#).

During the program, participants increased the length and intensity of their workouts over time. For example, participants received an individually tailored, progressive walking routine, and they wore a pedometer and recorded their daily steps over six weeks, starting on their first day of chemotherapy treatment. They were encouraged to gradually increase their steps by five to 20 percent every week. For resistance exercise, they performed exercises with therapeutic exercise bands. Participants were given individually-tailored workout plans that encouraged them to perform 10 required exercises (such as squats and chest presses) and four optional exercises daily. Participants were also encouraged to increase the intensity and number of repetitions of resistance band exercises gradually over the course of the program.

The researchers concluded that a low- to moderate-intensity home-based exercise program improved anxiety, mood, and social and emotional well-being for older patients with cancer who received chemotherapy treatments.

The researchers also noted that in the study, the people who benefited the most from the [exercise](#) program were [older adults](#) who received

chemotherapy and started off with worse anxiety, mood, and social and emotional well-being.

More information: Kah Poh Loh et al, Effects of a Home-based Exercise Program on Anxiety and Mood Disturbances in Older Adults with Cancer Receiving Chemotherapy, *Journal of the American Geriatrics Society* (2019). DOI: [10.1111/jgs.15951](https://doi.org/10.1111/jgs.15951)

Provided by American Geriatrics Society

Citation: Exercise improves anxiety and mood in older adults undergoing chemotherapy (2019, July 3) retrieved 4 May 2024 from <https://medicalxpress.com/news/2019-07-anxiety-mood-older-adults-chemotherapy.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--