

# Some children are more likely to suffer depression long after being bullied

July 1 2019

---



Credit: CC0 Public Domain

Some young adults who were bullied as a child could have a greater risk of ongoing depression due to a mix of genetic and environmental factors according to a new study from the University of Bristol.

Researchers wanted to find out what factors influenced [depression](#) in young adults between the ages of 10 and 24 and why some people responded differently to risk factors such as bullying, maternal postnatal depression, early childhood anxiety and domestic violence.

Using detailed mood and feelings questionnaires and [genetic information](#) from 3,325 teenagers who are part of Bristol's Children of the 90s study, alongside evidence of these risk factors at nine points in time they found that childhood bullying was strongly associated with trajectories of depression that rise at an early age. Children who continued to show high depression into adulthood were also more likely to have genetic liability for depression and a mother with postnatal depression. However, Children who were bullied but did not have any genetic liability for depression showed much lower depressive symptoms as they become young adults.

University of Bristol Ph.D. student Alex Kwong commented:

"Although we know that depression can strike first during the [teenage years](#) we didn't know how risk factors influenced change over time. Thanks to the Children of the 90s study, we were able to examine at multiple time points the relationships between the strongest risk factors such as bullying and maternal depression, as well as factors such as genetic liability.

"It's important that we know if some children are more at risk of depression long after any childhood bullying has occurred. Our study found that [young adults](#) who were bullied as children were eight times more likely to experience depression that was limited to childhood. However, some children who were bullied showed greater patterns of depression that continued into adulthood and this group of children also showed genetic liability and family risk.

"However, just because an individual has genetic liability to depression does not mean they are destined to go on and have depression. There are a number of complex pathways that we still don't fully understand and need to investigate further.

"The next steps should continue to look at both genetic and environmental risk factors to help untangle this complex relationship that would eventually help influence prevention and coping strategies for our health and education services."

Lecturer in Psychiatric Epidemiology at the University of Bristol Dr. Rebecca Pearson added:

"The results can help us to identify which groups of children are most likely to suffer ongoing symptoms of depression into adulthood and which children will recover across adolescence. For example, the results suggest that children with multiple risk factors (including [family history](#) and bullying) should be targeted for early intervention but that when [risk factors](#) such as bullying occur in isolation, symptoms of depression may be less likely to persist."

**More information:** Alex S. F. Kwong et al. Genetic and Environmental Risk Factors Associated With Trajectories of Depression Symptoms From Adolescence to Young Adulthood, *JAMA Network Open* (2019). [DOI: 10.1001/jamanetworkopen.2019.6587](https://doi.org/10.1001/jamanetworkopen.2019.6587)

Provided by University of Bristol

Citation: Some children are more likely to suffer depression long after being bullied (2019, July 1) retrieved 1 May 2024 from <https://medicalxpress.com/news/2019-07-children-depression-bullied.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.