


Deep brain stimulation modifies memory

July 29 2019



The location of posterior cingulate cortex, the stimulated brain region. Credit: Natu et al., *JNeurosci* 2019

Deep brain stimulation of the cingulate cortex worsens memory recall, according to research in epilepsy patients published in *JNeurosci*. The technique could be a way to target specific brain areas in the treatment of memory disorders.

Vaidehi Natu and colleagues at the University of Texas Southwestern

Medical Center stimulated the [posterior cingulate cortex](#), a brain region involved in memory, using electrodes inserted into the brains of epilepsy patients to treat seizures. The participants were shown a list of words, distracted, and then asked to recall the words.

Natu's team discovered that stimulating the cingulate cortex reduced the number of recalled words and modified hippocampal brain waves compared to no stimulation. This establishes a direct role for the [cingulate cortex](#) in memory encoding and demonstrates that memories can be manipulated via deep brain stimulation, which has implications for the treatment of memory disorders.

More information: *JNeurosci* (2019). [DOI: 10.1523/JNEUROSCI.0698-19.2019](#).
www.biorxiv.org/content/10.1101/497818v1

Provided by Society for Neuroscience

Citation: Deep brain stimulation modifies memory (2019, July 29) retrieved 5 May 2024 from <https://medicalxpress.com/news/2019-07-deep-brain-memory.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
