

Depression is common and linked with early death in patients with blood cancers

July 3 2019



Credit: CC0 Public Domain

In a *Psycho-Oncology* study of patients newly diagnosed with lymphoma or multiple myeloma, one-third of participants reported depressive symptoms around the time of diagnosis, and depressive symptoms were

linked with shorter survival. Shorter survival was observed in both patients who remained depressed and patients who recovered from depressive symptoms.

The study included 255 [patients](#), of whom 83 had depression. Over a median follow-up of approximately 3.5 years, 61 patients died. These included 30 patients in the "never depression" group, 6 patients in the "new-onset" group, 16 patients in the "remission" group, and 9 patients in the "persistent" group. These groups were based on [depressive symptoms](#) assessed before starting [cancer treatment](#) and one month later.

The risk of death was 3-times and 2-times higher in the "remission" and "persistent" groups, respectively, than in the "never depression" group; however, risk of death was not significantly greater in the "new-onset" group compared with the "never depression" group.

More information: Takaaki Hasegawa et al, Depressive symptoms during the first month of chemotherapy and survival in patients with hematological malignancies: A prospective cohort study, *Psycho-Oncology* (2019). [DOI: 10.1002/pon.5143](https://doi.org/10.1002/pon.5143)

Provided by Wiley

Citation: Depression is common and linked with early death in patients with blood cancers (2019, July 3) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-07-depression-common-linked-early-death.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--