

Discovery in mice points to potential treatment for vestibular disorders

July 9 2019



Credit: CC0 Public Domain

Researchers at the Stanford University School of Medicine have found a way to regenerate hair cells in the inner ears of mice, allowing the animals to recover vestibular function. It's the first time such recovery has been observed in mature mammals.

If further research shows that the technique can be applied to humans, it

would be an initial step toward treating vestibular disorders, such as dizziness. There is currently no [effective treatment](#) for dizziness and balance disorders caused by damaged or lost vestibular [hair cells](#). The only available therapy is teaching patients coping mechanisms through [physical therapy](#).

"This disabling condition is very common among the elderly, and one of the primary causes of falls," said Alan Cheng, MD, associate professor of otolaryngology-head and [neck surgery](#).

Cheng is the senior author of a paper about the research, which will be published July 9 in *Cell Reports*. Zahra Sayyid, an MD-Ph.D. student at Stanford, is the lead author.

The [hair cells](#) in the utricle, a section of the inner ear, help maintain balance and spatial orientation and regulate eye movement. Some antibiotics can damage these cells. Damage can also occur from infections or genetic disorders, or as a result of aging. In mature mammals, vestibular hair cells regenerate on their own only minimally. (Birds and fish, however, have the ability to completely regenerate them.)

In the United States, about 69 million people experience vestibular dysfunction, some because of problems with inner ear hair cells. They can feel as if they're spinning, lose their balance easily, suffer from nausea and have trouble tracking objects with their eyes. The symptoms can prevent patients from engaging in activities, including exercise and driving, and can lead to [social isolation](#).

To study these vestibular disorders, the researchers impaired the inner ear hair cells of mice and measured how well they regenerated on their own. The researchers found that about a third of the cells regenerated spontaneously but appeared immature, and vestibular function was

inconsistent.

Next, they manipulated Atoh1, a transcription factor that regulates hair cell formation, in the mice. In the animals that overexpressed Atoh1, as much as 70% of hair cells regenerated. The regenerated cells appeared relatively mature, and about 70% of these mice recovered vestibular function.

"This is very exciting. It's an important first step to find treatment for vestibular disorders," Cheng said. "We couldn't get sufficient regeneration to recover function before."

The researchers plan to study how other methods to enhance Atoh1's function may affect regeneration.

While the finding is a proof of concept, "it has opened the door for many more possibilities that could lead to treatment in people with vestibular [disorders](#)," Sayyid said.

Provided by Stanford University Medical Center

Citation: Discovery in mice points to potential treatment for vestibular disorders (2019, July 9) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2019-07-discovery-mice-potential-treatment-vestibular.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--