

Early EEG helps predict cardiac arrest outcomes in comatose

July 17 2019



(HealthDay)—Early electroencephalography (EEG) reliably predicts the

outcome of comatose patients after cardiac arrest, according to a study recently published in the *Annals of Neurology*.

Barry J. Ruijter, M.D., Ph.D., from the University of Twente in Enschede, Netherlands, and colleagues assessed whether early EEG can reliably predict outcomes among comatose patients after cardiac arrest. Blinded reviewers assessed five-minute EEG epochs at eight predefined time points from six hours to five days after cardiac arrest.

The researchers found good outcomes for 46 percent of 850 patients. At six or more hours after cardiac arrest, generalized suppression and synchronous patterns with ≥ 50 percent suppression predicted poor outcome without [false positives](#). At 12 and 24 hours after cardiac arrest, the summed sensitivity was 0.47 (95 percent confidence interval [CI], 0.42 to 0.51) and 0.30 (95 percent CI, 0.26 to 0.33), respectively, with a specificity of 1.00 at both time points (95 percent CI, 0.99 to 1.00). At ≥ 36 hours, sensitivity for poor outcome was ≤ 0.22 . In contrast, continuous EEG patterns at 12 hours predicted good outcome, with a sensitivity of 0.50 (95 percent CI, 0.46 to 0.55) and specificity of 0.91 (95 percent CI, 0.88 to 0.93), while specificity for the prediction of good outcome at 24 hours or later was

Citation: Early EEG helps predict cardiac arrest outcomes in comatose (2019, July 17) retrieved 26 June 2024 from <https://medicalxpress.com/news/2019-07-early-eeeg-cardiac-outcomes-comatose.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.