

Helicopter transport for stroke patients decreases time to surgery, new study finds

July 22 2019

The sooner that a severe stroke patient can access thrombectomy, the more likely they are to experience a good outcome. A new study shows that using emergency helicopter ambulance services to transfer a patient to a hospital that can perform a stroke thrombectomy—a minimally invasive surgery which removes the blood clot in the brain causing the stroke—ensures faster access to potentially life-saving care.

The study—An Analysis of Stroke Thrombectomy Interhospital Transportation Modality—took place between January 2015 and March 2018 and was released today at the 16th Annual Meeting of the Society of NeuroInterventional Surgery. The researchers behind the study aimed to determine if ground or air ambulance transportation methods allowed a patient to access thrombectomy the fastest.

The researchers analyzed 133 patients who were transferred to Rush University Medical Center in Chicago, a comprehensive stroke center, for thrombectomy. Patients who were transported to Rush University by helicopter had a significantly shorter time to [surgery](#) start. Furthermore, air transport reduced time more as distances between hospitals increased, getting patients to surgery on average 42 minutes faster when transported more than 30 miles. However, all transport of less than 10 miles was done by ground ambulance.

"We know that when a patient can receive earlier appropriate stroke care, they experience better outcomes. We should consider this study as evidence that transferring a patient by helicopter ambulance to a hospital

that can perform a thrombectomy will likely provide the patient the best chance of recovering from a major stroke," says Dr. Hormuzdiyar Dasenbrock, first author of the study from Rush University Medical Center. "However, [patient safety](#) is the first concern, and sometimes air transport cannot be used due to weather; also, [air transport](#) may not be faster if the distance between hospitals is less than 10 miles."

This research reinforces a recent report from the Centers for Disease Control and Prevention's Division for Heart Disease and Stroke Prevention, which found that patients experienced decreased time and access to receiving crucial [stroke](#) medications (intravenous thrombolytics) when they were transported via air to an appropriate facility, and extends these findings to patients undergoing [thrombectomy](#)

Provided by Society of NeuroInterventional Surgery

Citation: Helicopter transport for stroke patients decreases time to surgery, new study finds (2019, July 22) retrieved 25 April 2024 from <https://medicalxpress.com/news/2019-07-helicopter-patients-decreases-surgery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--