

# High rates of opioid prescriptions for osteoarthritis

July 8 2019

---



Credit: CC0 Public Domain

Opioids work against severe pain but the risks of side effects and addiction are high. In the U.S. alone, 26 people die every day from overdoses. Now researchers in an international collaboration have



investigated how common opioid prescriptions are for osteoarthritis patients in Sweden. It emerged that every fourth patient was prescribed opioids at some point between November 2014 and October 2015.

"The results show an alarmingly high rate of [opioid](#) prescriptions for [osteoarthritis patients](#)," says Martin Englund, professor at Lund University and physician at Skåne University Hospital who led the study in collaboration with researchers from Denmark and the United Kingdom.

The researchers based their research on the population registry in southern Sweden of people who were at least 35 years old (a total of 751 579 people) and who had received a diagnosis of knee or hip osteoarthritis at some point during 1998-2015 (a total of 82 379 people). With the help of data from the Swedish Prescribed Drug Register the researchers were able to see how many of these people were prescribed opioid preparations between 1 November 2014 and 31 October 2015. This was reported to be every fourth osteoarthritis patient (24 percent).

The study also estimated that at least 12 percent of all new opioid prescriptions are given to patients with osteoarthritis. The most frequently prescribed drugs were codeine, tramadol, oxycodone and morphine preparations.

"The result is remarkable. Opioids are highly addictive preparations and are not on the list of recommended treatments for osteoarthritis in Sweden, except for in extreme cases or in conjunction with surgical treatment. However, even when we removed all prescription in conjunction with surgery, the result was very similar," says Martin Englund.

In Sweden there is a tradition of collecting [personal data](#) in a register that is connected to the individual's own personal identification number.



Such data is not typically available in most countries.

"In general, a high use, or an increase in the use of opioids among osteoarthritis patients have been reported in the US, Catalonia in Spain and in Australia. But to estimate reliable figures of opioid prescriptions preferably requires careful documentation in registers covering the entire population, and from all levels of health care," says Martin Englund.

He says that the results indicate a significant challenge within the field of osteoarthritis: there is good basic treatment for patients—osteoarthritis education and exercise programs—however, many osteoarthritis patients do not utilize this. Subsequently, or if there is insufficient improvement after basic treatment, there is a significant lack of good pharmaceutical non-surgical treatment alternatives to offer the patient, other than some relief of symptoms.

"If the patient has developed severe osteoarthritis, and other treatments do not offer sufficient relief of symptoms, it is possible to offer prosthetic surgery. However, it is substantial and expensive [surgical treatment](#), and should only be offered to patients that really need the operation. Unfortunately the healthcare system is lacking in pharmaceutical [treatment](#) options that can relieve pain and modify the development of the disease itself, i.e. have an impact on the actual biological process. We need more research on this," says Martin Englund.

**More information:** J.B. Thorlund et al. Opioid use in knee or hip osteoarthritis: a region-wide population-based cohort study, *Osteoarthritis and Cartilage* (2019). [DOI: 10.1016/j.joca.2019.01.005](https://doi.org/10.1016/j.joca.2019.01.005)



Provided by Lund University

Citation: High rates of opioid prescriptions for osteoarthritis (2019, July 8) retrieved 5 May 2024 from <https://medicalxpress.com/news/2019-07-high-opioid-prescriptions-osteoarthritis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.