

Study links air pollution to increase in newborn intensive care admissions

July 19 2019

NIH study links air pollution to increase in newborn intensive care admissions

Infants born to women exposed to [high levels of air pollution](#) in the week before delivery are more likely to be admitted to a newborn [intensive care unit](#) (NICU), suggests an analysis by researchers at the National Institutes of Health. Depending on the type of [pollution](#), chances for NICU [admission](#) increased from about 4% to as much as 147%, compared to infants whose mothers did not encounter high levels of air pollution during the week before delivery. The study was led by Pauline Mendola, Ph.D., of the Epidemiology Branch at NIH's Eunice Kennedy Shriver National Institute of Child Health and Human Development. It appears in *Annals of Epidemiology*.

"Short-term exposure to most types of air pollutants may increase the risk for NICU admission," Dr. Mendola said. "If our findings are confirmed, they suggest that pregnant women may want to consider limiting their time outdoors when air quality advisories indicate unhealthy conditions."

Previous studies have linked elevated levels of certain kinds of air pollutants to higher risks for gestational diabetes and preeclampsia, a blood pressure disorder of pregnancy. Earlier research also has shown that infants born to women exposed to high levels of air pollutants are at risk for [preterm birth](#), of being small for their gestational age at birth and of growing more slowly than normal in the uterus. Given these

associations, the study authors sought to determine whether prenatal exposure to air pollution might increase the chance for NICU admission.

Researchers analyzed data from the Consortium on Safe Labor, which compiled information on more than 223,000 births at 12 clinical sites in the United States from 2002 to 2008. They linked records from more than 27,000 NICU admissions to data modified from the Community Multiscale Air Quality Modeling System, which estimates environmental pollution concentrations in the United States. Researchers matched air quality data in the area where each birth occurred to the week before delivery, the day before delivery and the day of delivery. They then compared these time intervals to air quality data two weeks before delivery and two weeks after delivery to identify risk of NICU admission associated with pollution levels.

The researchers also examined the odds of NICU admission associated with high concentrations of particulate matter (pollution particles) less than 2.5 microns in diameter (PM_{2.5}). These types of particles originate from various sources, among them diesel and gasoline engines, power plants, landfills, sewage facilities and industrial processes. Exposure to high concentrations of organic compounds in the air was associated with a 147% increase in risk of NICU admission. Elemental carbon and ammonium ions presented similar increases in risk (38% and 39%, respectively), while exposure to nitrate compounds was associated with a 16% higher risk of NICU admission.

Chances of NICU admission increased significantly with exposures to traffic-related pollutants on the day before and the day of delivery, compared to the week before delivery: 4% and 3%, respectively, for an approximately 300 parts per million (ppm) increase in carbon monoxide; 13% and 9% for an approximately 26 ppm increase in [nitrogen dioxide](#); and 6% and 3% an approximately 3 ppm increase in sulfur dioxide.

Researchers do not know why exposure to air pollution might increase the chances for NICU admission. They theorize, however, that pollutants increase inflammation, leading to impaired blood vessel growth, particularly in the placenta, which supplies oxygen and nutrients to the developing fetus.

The authors note that rising NICU admission rates present financial challenges for families and society, as average daily NICU costs may reach or exceed \$3,000. If their results are confirmed by other studies, limiting [pregnant women](#)'s exposure to high levels of air pollutants may provide a way to reduce NICU admissions.

More information: Indulaxmi Seeni et al, Acute air pollution exposure and NICU admission: a case-crossover analysis, *Annals of Epidemiology* (2019). [DOI: 10.1016/j.annepidem.2019.07.008](https://doi.org/10.1016/j.annepidem.2019.07.008)

Provided by NIH/Eunice Kennedy Shriver National Institute of Child Health and Human Development

Citation: Study links air pollution to increase in newborn intensive care admissions (2019, July 19) retrieved 24 April 2024 from <https://medicalxpress.com/news/2019-07-links-air-pollution-newborn-intensive.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--