

## Sex differences in kidney injury

July 29 2019, by Kelsey Herbers



Credit: CC0 Public Domain

Evidence indicates men have higher incidence of chronic kidney disease and quicker renal decline than premenopausal women. However, sex differences in the expression and activation of the EGF receptor (EGFR), which mediates progressive renal injury, are largely

![](_page_1_Picture_0.jpeg)

unexplored.

Ming-Zhi Zhang, MD, Raymond Harris, MD, and colleagues examined EGFR expression in mice, human kidney tissue and cultured cells. The results, published in *Journal of the American Society of Nephrology*, indicated lower EGFR levels in females than males in mice and adult human kidneys.

In mice with an activated EGFR, males showed multiple types of renal injury while females showed minimal injury. Castration decreased EGFR expression in males and protected against kidney injury, while testosterone increased EGFR expression and renal injury in females.

The results indicate <u>sex differences</u> in susceptibility to progressive <u>kidney</u> injury—which may be mediated by testosterone—are in part due to differences in renal EGFR expression.

**More information:** Ming-Zhi Zhang et al. The Role of the EGF Receptor in Sex Differences in Kidney Injury, *Journal of the American Society of Nephrology* (2019). DOI: 10.1681/ASN.2018121244

Provided by Vanderbilt University

Citation: Sex differences in kidney injury (2019, July 29) retrieved 21 May 2024 from https://medicalxpress.com/news/2019-07-sex-differences-kidney-injury.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.