

Uranium toxicity may be causing high rates of obesity and diabetes in Kuwait

July 8 2019



Dr. Max Goodson Credit: Dr. Max Goodson

Kuwait has some of the highest rates of obesity and diabetes in the world, and scientists don't know why. This question was addressed by Dr. Max Goodson, Emeritus Professor at the Forsyth Institute.

Goodson is the principal investigator of a large-scale study that analyzed diet, salivary bacteria, salivary protein biomarkers, and salivary metabolites of 10-year-old Kuwaiti children. Researchers collected saliva samples from 94 children once in 2012 and again in 2014.

All of the children were healthy at the start of the study, but approximately half of them developed obesity and metabolic syndrome two years later. When analyzing differences in metabolites found in saliva samples, Goodson and his colleagues discovered elevated levels of a biomarker for uranium toxicity were present in the saliva of children who became obese. The results were published recently in *Frontiers in Endocrinology*.

The biomarker, a compound called N1-Methyl-2-pyridone-5-carboxamide (2PY), is a metabolite of the vitamin niacin. It is present at low levels in almost everyone's [saliva](#). However, 2PY was found at higher than normal concentrations only in the [saliva samples](#) of the obese children in Kuwait. In a comparable group of obese children from Maine and Massachusetts, elevated levels of 2PY in obese children were not found.

"The implication," Goodson said, "is that these [children](#) may be suffering from uranium toxicity, which may be contributing to the high rates of obesity and diabetes in Kuwait."

The idea that uranium toxicity would be a health concern in Kuwait makes sense. In 1990-1991, the United States dropped almost 300 tons of uranium-containing munitions in the central region of the country, Goodson said. As part of the recent study, researchers found that the highest rates of obesity and salivary 2PY were concentrated in that central region and reduced in peripheral areas of Kuwait.

Goodson is careful to point out that the study measured a biomarker for

uranium, not uranium itself. More research is needed to determine whether toxicity from exposure to the [heavy metal](#) is truly the culprit for high rates of obesity and diabetes seen in Kuwait.

More information: Jo Max Goodson et al, Salivary N1-Methyl-2-Pyridone-5-Carboxamide, a Biomarker for Uranium Uptake, in Kuwaiti Children Exhibiting Exceptional Weight Gain, *Frontiers in Endocrinology* (2019). [DOI: 10.3389/fendo.2019.00382](https://doi.org/10.3389/fendo.2019.00382)

Provided by Forsyth Institute

Citation: Uranium toxicity may be causing high rates of obesity and diabetes in Kuwait (2019, July 8) retrieved 27 April 2024 from <https://medicalxpress.com/news/2019-07-uranium-toxicity-high-obesity-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.