

Anxiety, depression linked to more opioid use after surgery

August 22 2019



Credit: CC0 Public Domain

Surgeons wielding their life-saving scalpels, laparoscopic tools, or other implements to repair or remove what ails their patients understand all too well that pain is an unavoidable part of the healing process. Yet the

current opioid crisis has made the standard prescribing practices for these highly effective analgesics fraught with risk.

New research from Michigan Medicine could help clinicians mitigate that risk by identifying which patients are more likely to continue to use opioids after their immediate recovery period.

"There is not much research on which [surgical patients](#) require more or less opioids, despite a push in the field for personalized medicine," says first author Daniel Larach, M.D., M.T.R., M.A., a resident at U-M at the time of the study and now assistant professor of clinical anesthesiology at the University of Southern California.

"Often with [postoperative opioid](#) prescribing, personalization falls by the wayside, with surgeons using the same amounts for every person receiving a certain procedure."

He and the study team looked at data for more than 1,000 people undergoing an elective hysterectomy, [thoracic surgery](#), or a total knee or hip replacement. Before their procedures, each patient provided demographic information and filled out several screening questionnaires. They were given scores measuring their degree of depression, anxiety, fatigue, sleep disturbance, physical function, as well as the severity of their overall and surgical site pain. The research team also measured how many pills were prescribed per patient. Each patient was then contacted one month following surgery to assess how many [opioid](#) pills they had consumed.

"We found that anxiety is linked with more opioid use, which is disheartening to see but also heartening in the sense that this is something we could potentially target," says Larach.

Other patient factors linked to increased [opioid use](#) included younger

age, non-white race, no college degree, alcohol and tobacco use, and sleep disturbance.

Chad Brummett, M.D., associate professor of anesthesiology and director of anesthesia [clinical research](#) and pain research, says people may be knowingly or unknowingly medicating for other conditions.

"The only thing we're giving them is opioids and we're not giving them alternatives or other options," he says. For example, patients with high anxiety around the time of surgery could be offered behavioral care or other non-opioid medications for anxiety and resulting pain.

Brummett, senior author on the *Annals of Surgery* paper, also notes that this study found overprescription of opioids for all surgical procedures and a correlation between the prescription size and use.

"I think it is striking that you see once again that the more you prescribe, the more patients take, even after adjusting for all of these other risk factors," explains Brummett.

Larach and Brummett note that right-sizing prescriptions through initiatives such as the Michigan Opioid Prescribing Engagement Network (OPEN), which provides recommendations for prescription amounts for various medical procedures, is a critical first step. But, they say, this step should be followed by more research into specific patient factors that can be addressed in other ways.

Says Brummett, "We are asking surgeons to learn about and think about pain and behavioral health in ways that we have not previously done. It will require an open mind."

More information: Daniel B. Larach et al, Patient Factors Associated with Opioid Consumption in the Month Following Major Surgery,

Annals of Surgery (2019). [DOI: 10.1097/SLA.0000000000003509](https://doi.org/10.1097/SLA.0000000000003509)

Provided by University of Michigan

Citation: Anxiety, depression linked to more opioid use after surgery (2019, August 22) retrieved 24 April 2024 from

<https://medicalxpress.com/news/2019-08-anxiety-depression-linked-opioid-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.