

Cancer survivors likely to face increase in long-term risk of cardiovascular disease

August 20 2019

In one of the largest studies of its kind, the research team, led by the London School of Hygiene & Tropical Medicine, analysed the medical records of more than 630,000 people in the UK, including over 100,000 survivors of a range of cancers.

They found raised risks of blood clots forming in the veins among survivors of most cancers, ranging from approximately a two- to ten-fold increase in risk for different [cancer](#) sites. Risks decreased over time but were still elevated for more than five years after the cancer diagnosis. Survivors of half of the 20 cancers studied also faced increased risks of heart muscle damage, known as cardiomyopathy, and [heart failure](#).

For example, survivors of blood, oesophagus, lung, kidney and [ovarian cancers](#) were more than one and a half times more likely to experience cardiomyopathy or heart failure than people with no prior cancer.

An increased risk of coronary artery disease and stroke was found in some cancer survivors, including those with prior blood cancers.

The researchers stress that for many people the overall risks will still be low, particularly for younger cancer survivors, but say the findings highlight the need for new strategies to prevent and manage heart and circulation problems in cancer survivors, including a need to raise awareness among GPs of the raised risks.

While the researchers could not definitively identify the causes of the

increased risks, the team's analysis suggests that exposure to cancer treatments such as chemotherapy is likely to play a key role.

Lead author Helen Strongman from the London School of Hygiene & Tropical Medicine said: "Over recent decades [cancer treatment](#) and management have improved substantially, resulting in a large and growing population of long-term cancer survivors. Around half of those diagnosed with cancer in developed countries are now expected to survive for more than 10 years. However, there are concerns that there may be increased long-term risks of cardiovascular disease following [cancer diagnosis](#), driven by treatment side-effect effects and the potential impact of the cancer itself."

While previous studies have demonstrated the short- to medium-term increased risk of cardiovascular disease from some specific cancer treatments, there have been limited data on the overall and long-term differences in cardiovascular risk between cancer survivors and those who have never had cancer.

To address this, the team brought together anonymised data from primary care, hospitals, cancer registries, and death certificates, to quantify the absolute and relative risks of a comprehensive range of cardiovascular diseases in survivors of the 20 most common adult cancers compared with cancer-free general population controls. The research also accounted for other risk factors for cardiovascular disease, such as older age, smoking and body mass index.

The team found large increases in risk of venous thromboembolism (blood clots in the veins) among survivors of 18 of the 20 cancers studied compared to cancer-free controls. Survivors of breast cancer had double the risk of venous blood clots one year after diagnosis compared with similar women with no prior cancer, though the overall level of risk remained low, especially for younger survivors.

For female breast cancer survivors aged under 60 years, six extra blood clot cases per year were observed per 1,000 women, while among older women (80+ years) 12 extra cases per year were observed per 1,000 women. These risks decreased over time but remained elevated for at least five years after diagnosis with most cancers.

The team also observed long-term increased risks of heart failure and heart muscle damage among survivors of 10 of the 20 site-specific cancers studied. Among people who had non-Hodgkin lymphoma five years previously, there were four extra cases of heart failure per year for every 1,000 people aged under 60, rising to 21 extra cases per year for every 1000 people aged over 80 years.

The senior author of the study Professor Krishnan Bhaskaran from the London School of Hygiene & Tropical Medicine said: "With treatment for cancer becoming more effective, we must start thinking about living beyond cancer and improving the health of survivors. Our work revealed raised risks of various cardiovascular diseases in cancer survivors. These findings are important because those affected might benefit from preventative measures and earlier interventions to reduce the impact.

"If patients, primary care doctors, and specialists are more aware of the potential for increased cardiovascular risks in this patient group, we might detect problems earlier and improve outcomes. There is limited guidance at present to help doctors manage cardiovascular risk in cancer survivorship, and further evidence is needed on whether routine monitoring or additional preventative measures would benefit those with the highest risks."

The authors acknowledge limitations of their study including that there was only limited information available about the anti-cancer treatments that patients had received. Although the research team had some information on whether patients had received chemotherapy,

radiotherapy, and surgery, there was no detailed information on specific chemotherapy drugs, radiotherapy doses, or surgery procedures.

More information: *The Lancet*, www.thelancet.com/journals/lan...
[http://www.thelancet.com/journals/lan... \(19\)31674-5/fulltext](http://www.thelancet.com/journals/lan...)

Provided by London School of Hygiene & Tropical Medicine

Citation: Cancer survivors likely to face increase in long-term risk of cardiovascular disease (2019, August 20) retrieved 25 April 2024 from <https://medicalxpress.com/news/2019-08-cancer-survivors-long-term-cardiovascular-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.