

Diabetes treatment targets have not improved in the US since 2005

August 12 2019



Credit: CC0 Public Domain

More than 30 million Americans are living with diabetes. Diabetes treatment is generally focused on controlling blood sugar, blood pressure and cholesterol level, as well as promoting smoking cessation. A new study by investigators at Massachusetts General Hospital (MGH) published online in *JAMA Internal Medicine* suggests that the

achievement of these targets has not improved for U.S. adults with diabetes since 2005.

Using 2005-2016 nationally representative survey and examination data from Centers for Disease Control and Prevention (CDC), the authors conducted statistical analyses to evaluate the cascade of diabetes care in the United States which is defined as diabetes diagnosis, linkage to care and achievement of treatment targets. They found that more than one in four U.S. adults with diabetes were not diagnosed and nearly one in three were not receiving appropriate care for diabetes in the study period.

"Fewer than one in four American adults with diagnosed diabetes achieve a controlled level of [blood sugar](#), [blood pressure](#) and cholesterol and do not smoke tobacco. Our results suggest that, despite major advances in diabetes drug discovery and movement to develop innovative care delivery models over the past two decades, achievement of diabetes care targets has not improved in the United States since 2005," says lead and corresponding author Pooyan Kazemian, Ph.D., of the MGH Medical Practice Evaluation Center, and instructor in Medicine at Harvard Medical School (HMS).

The authors also investigated potential age, gender and racial disparities in U.S. diabetes care. They found that certain populations demonstrated persistent disparities in meeting diabetes care targets over the study period.

"Younger age (18-44), female and non-white adults with diabetes had lower odds of achieving the composite [blood sugar](#), blood pressure, cholesterol and non-smoking target," says Kazemian.

Insurance coverage was the strongest indicator of diagnosis, linkage and achievement of the composite treatment targets.

"Barriers accessing [health care](#), including lack of health insurance and high drug costs, remain major factors that have not been adequately addressed on a population level," says senior author Deborah J. Wexler, MD, MSc, of the MGH Diabetes Unit, and associate professor in Medicine at HMS.

The authors note that continued monitoring of diabetes care cascade can help shed light on the population-level steps required to improve diabetes care in the United States.

"Treatment advances in diabetes mellitus can meaningfully improve outcomes only if they effectively reach the populations at risk. Our findings suggest this is not the case in the U.S. and indicate an immediate need for better approaches to [diabetes](#) care delivery including a continued focus on reaching underserved populations with persistent disparities in care," says Wexler.

More information: *JAMA Internal Medicine* (2019). [DOI: 10.1001/jamainternmed.2019.2396](#)

Provided by Massachusetts General Hospital

Citation: Diabetes treatment targets have not improved in the US since 2005 (2019, August 12) retrieved 23 June 2024 from <https://medicalxpress.com/news/2019-08-diabetes-treatment.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.