

Healthier homes could cut hospital stays for young children, researchers find

August 15 2019



Credit: CC0 Public Domain

Almost 20 percent of hospital admissions of young children with acute respiratory infections could be prevented if their houses were free from damp and mold, researchers have found.

The research, published in the international medical journal *Thorax* today, is the largest ever case-control study of the association between housing quality and hospital admissions for young children with [acute respiratory infections](#).

Lead author, Dr. Tristram Ingham, a Senior Research Fellow at the University of Otago, Wellington, says reducing exposure to poor quality housing could cut the number of hospital admissions for the under-twos by 1,700 (19 percent) a year.

In 2015, 9,003 under-two-year-olds were hospitalized in Wellington for acute respiratory infections out of an estimated population of 118,580, a rate per population of 7.6 percent.

"If all housing was free from damp and mold, the country could save just under \$8 million a year solely in hospital costs," Dr. Ingham says.

"Improving housing quality to prevent such hospitalizations is also highly desirable in terms of future costs, with poor [housing quality](#) in childhood associated with adverse health outcomes in adulthood."

The researchers conducted the case-control study in two pediatric wards and five general practice clinics in Wellington during winter and spring seasons between 2011 and 2013. The quality of housing was assessed using the Respiratory Hazard Index (RHI), a 13-point scale of household quality factors including an eight-item damp/mold subscale.

The researchers studied 188 children who were admitted to hospital with acute respiratory infections and 454 control patients—those who saw their GP either with a respiratory illness which did not require hospitalization or for a routine immunization.

The children who became ill had significantly higher levels of dampness

and mold in their homes, which could be addressed by a number of measures, including better heating, insulation and ventilation.

The research was part of a wider Whiti Te Rā Case-Control Study which examines the broader risk factors for hospitalization for acute respiratory [infection](#) using a kaupapa Māori research framework.

Acute respiratory infections are a major cause of early childhood illness, and account for 50 percent of consultations general practitioners have with children under five. Upper respiratory tract infections are the most common, but acute respiratory infections also include [lower respiratory tract infections](#), such as bronchitis and bronchopneumonia.

The [research paper](#), "Damp moldy [housing](#) and early childhood hospital admissions for acute respiratory infection: a case control study," is published in *Thorax*.

More information: Tristram Ingham et al. Damp mouldy housing and early childhood hospital admissions for acute respiratory infection: a case control study, *Thorax* (2019). [DOI: 10.1136/thoraxjnl-2018-212979](https://doi.org/10.1136/thoraxjnl-2018-212979)

Provided by University of Otago

Citation: Healthier homes could cut hospital stays for young children, researchers find (2019, August 15) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-08-healthier-homes-hospital-young-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.