

High fat foods can increase CBD absorption into the body

August 13 2019

While oral cannabidiol (CBD) capsules were approved by the U.S. Food and Drug Administration (FDA) for use in patients with seizures in 2018, very little was known about the effect of food on CBD absorption.

A University of Minnesota study, published in *Epilepsia*, examined whether eating high-fat foods after taking CBD increased the body's [absorption](#) of CBD. The study tested whether fasting or a high fat meal has an effect when cannabidiol oral capsules were taken by patients.

To find out what effect a fatty meal would have on CBD absorption, the research group measured CBD concentrations in epilepsy patients at the MINCEP Epilepsy Care clinic who were taking 99 percent pure CBD capsules. Concentrations from patients who took CBD on an empty stomach and a standardized fatty breakfast (i.e. breakfast burrito) were compared.

"The type of [food](#) can make a large difference in the amount of CBD that gets absorbed into the body. Although fatty foods can increase the absorption of CBD, it can also increase the variability as not all meals contain the same amount of fat," said Angela Birnbaum, a professor in the College of Pharmacy and study co-author.

"Increases in the amount of the CBD dose being absorbed into the body can also lead to lower medication costs," said Ilo Leppik, study co-author, a professor in the College of Pharmacy and an adjunct professor at the Medical School.

The study found:

- CBD exposure is vastly increased when CBD is taken with high fatty foods;
- when compared to fasting, taking CBD with food increased the amount of CBD in the body by four-times and the maximum amount recorded in the participants' blood by 14-times;
- no cognitive differences were identified, which is consistent with previous studies.

"For epilepsy patients, a goal is to maintain consistent blood concentrations of drug," said Birnbaum. "This study shows that CBD concentrations could vary significantly if patients take it differently, sometimes with or without food. Variations in blood concentrations could leave a patient more susceptible to seizures."

More information: Angela K. Birnbaum et al, Food effect on pharmacokinetics of cannabidiol oral capsules in adult patients with refractory epilepsy, *Epilepsia* (2019). [DOI: 10.1111/epi.16093](https://doi.org/10.1111/epi.16093)

Provided by University of Minnesota

Citation: High fat foods can increase CBD absorption into the body (2019, August 13) retrieved 6 July 2024 from <https://medicalxpress.com/news/2019-08-high-fat-foods-cbd-absorption.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.