

Individualised approach to identify 'fertile windows' could benefit many women

August 26 2019



Credit: CC0 Public Domain

Menstrual cycles are considerably varied with only 13% of women having cycles that last 28 days, according to a new study led by UCL and Natural Cycles, a contraceptive app.



The study, published today in *npj Digital Medicine*, is one of the biggest overviews of menstrual cycles to date and analysed data over 600,000 menstrual cycles and 124,648 women from Sweden, USA and the UK and could help couples better understand when pregnancy is likely and unlikely to occur.

Researchers set out to investigate menstrual cycle characteristics and associations with age, BMI and body temperatures. The findings show an average cycle length is 29.3 days and only around 13% of cycles are 28 days in length. Across the study, 65% of women had cycles that lasted between 25 and 30 days.

The menstrual cycle begins and ends with menstrual bleeding and is divided by ovulation into two phases: the follicular and luteal phases. Researchers found the average follicular phase length was 16.9 days and the average luteal phase length was 12.4 days. The average cycle length decreased by 0.18 days and average follicular phase length decreased by 0.19 days per year of age from 25 to 45 years. The average variation of cycle length for obese women was 0.4 days or 14% higher. Cycle length variability was observed to a lesser extent in non-obese and underweight women.

Co-author, Professor Joyce Harper (UCL Institute for Women's Health), said: "Our study is unique in analysing over half a million cycles and rewriting our understanding of the key stages. Traditionally studies have concentrated on women who have approximately 28 day cycles and these studies have formed our understanding of the menstrual cycle.

"For the first time our study shows that few women have the text book 28 day cycle, with some experiencing very short or very long cycles. We studied all women who used the app.

"We also demonstrate that ovulation does not occur consistently on day



14 and therefore it is important that women who wish to plan a pregnancy are having intercourse on their <u>fertile days</u>. In order to identify the fertile period, it is important to track other measures such as basal body temperature as cycle dates alone are not informative."

Co-author, Dr. Simon Rowland, Head of Medical Affairs at Natural Cycles, said: "Given the variations in cycle length and follicular phase length that we have described, especially for cycles outside the average range (25-30 days), an individualised approach to identify the fertile window should be adopted. Apps giving predictions of fertile days based solely on cycle dates could completely miss the fertile window and it is therefore unsurprising that several studies have shown that calendar apps are not accurate in identifying the fertile window.

"Besides the potential benefits to the individual, <u>fertility awareness apps</u> and the associated databases of fertility data provide a unique opportunity to examine a large number of menstrual cycles in order to improve understanding."

Data was collected from women using the app between September 2016 and February 2019 aged 18 to 45 and had a BMI between 15 and 50 and had not been using hormonal contraception within the 12 months prior to registration. Users who stated at registration that they had a pre-existing <u>medical condition</u> (polycystic ovarian syndrome, hypothyroidism or endometriosis) or who had menopausal symptoms were excluded.

The authors note the main limitation of this study is that the study population is derived solely from users of the app who may not be representative of the wider population. In particular, only 8% of women in our study were obese compared to 15% of <u>women</u> in the general population.

Professor Harper added: "The widespread use of mobile phone apps for



personal health monitoring is generating large amounts of data on the menstrual cycle. Provided that the real-world data can be validated against traditional clinical studies done in controlled settings, there is enormous potential to uncover new scientific discoveries.

"This is one of the largest ever analyses of <u>menstrual cycle</u> characteristics. These initial results only scratch the surface of what can be achieved. We hope to stimulate greater interest in this field of research for the benefit of public health."

More information: Jonathan R. Bull, Simon P. Rowland, Elina Berglund Scherwitzl, Raoul Scherwitzl, Kristina Gemzell Danielsson and Joyce Harper, 'Real-world menstrual cycle characteristics of more than 600,000 menstrual cycles' *Nature Digital Medicine* <u>doi.org/10.1038/s41746-019-0152-7</u>

Provided by University College London

Citation: Individualised approach to identify 'fertile windows' could benefit many women (2019, August 26) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2019-08-individualised-approach-fertile-windows-benefit.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.