


## Kids might be naturally immunized after C. difficile colonization in infancy

August 6 2019



This photograph depicts Clostridium difficile colonies after 48hrs growth on a blood agar plate; Magnified 4.8X. C. difficile, an anaerobic gram-positive rod, is the most frequently identified cause of antibiotic-associated diarrhea (AAD). It accounts for approximately 15–25% of all episodes of AAD. Credit: CDC

Exposure to *C. difficile* in infancy produces an immune response that might protect against this gastrointestinal infection later in childhood, according to a study published in *Clinical Infectious Diseases* journal. Researchers found that infants who were naturally exposed to *C. difficile* in the environment and became colonized with the bacteria had



antibodies in their blood. Analyses using a state-of-the-art assay revealed that these antibodies neutralized toxins that cause *C. difficile* infection, preventing harmful effects to cells exposed to these toxins. This suggests that a natural immunization occurs, although future studies will need to determine if it would prevent illness years later after another *C. difficile* exposure.

"We found an <u>immune response</u> in infants colonized with *C. difficile*, which might be beneficial as they get older, although we are still studying the extent and duration of this natural immunization," says lead author Larry Kociolek, MD, MSCI, from Stanley Manne Children's Research Institute at Ann & Robert H. Lurie Children's Hospital of Chicago, who is an Assistant Professor in Pediatrics-Infectious Diseases at Northwestern University Feinberg School of Medicine and Associate Medical Director of Infection Prevention and Control at Lurie Children's. "We are optimistic because we know from previous studies that adults with anti-toxin antibodies have a <u>lower risk</u> for illness from *C. difficile* <u>infection</u>."

*C. difficile* is a frequent cause of community- and healthcare-associated infection in adults and <u>children</u>. While roughly half of all infants get exposed, they normally do not get sick from these bacteria. Older children and adults usually get diarrhea that needs to be treated by antibiotics. A more severe form of the infection may cause inflammation of the colon that requires surgery and could be fatal. Children tend to have milder symptoms than adults. The pediatric incidence of *C. difficile* infection peaks in the 1-to-4-year age group and during teenage years.

"Given our results, we suspect that young children who get sick from *C*. *difficile* were probably not exposed as infants and so did not develop immunity," says Dr. Kociolek. "In adolescents, immunity might be waning. If with more research we can show that this is true, then there



might be a role for vaccinating susceptible children and teens against *C. difficile.*"

Currently, vaccines against *C. difficile* toxins are in clinical trials for adults. Pediatric clinical trials will be needed before a vaccine is available for children and adolescents.

**More information:** Larry K Kociolek et al, Natural Clostridioides difficile toxin immunization in colonized infants, *Clinical Infectious Diseases* (2019). DOI: 10.1093/cid/ciz582

Provided by Ann & Robert H. Lurie Children's Hospital of Chicago

Citation: Kids might be naturally immunized after C. difficile colonization in infancy (2019, August 6) retrieved 25 April 2024 from <a href="https://medicalxpress.com/news/2019-08-kids-naturally-immunized-difficile-colonization.html">https://medicalxpress.com/news/2019-08-kids-naturally-immunized-difficile-colonization.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.