

Psychiatric comorbidity contributes to increased mortality in ADHD

August 8 2019



Credit: CC0 Public Domain

Psychiatric comorbidity may play an important role in the increased risk of premature death in people with ADHD (attentiondeficit/hyperactivity disorder), according to a new extensive registry study conducted at Karolinska Institutet and Örebro University in Sweden. The results, which are published in JAMA Psychiatry, suggest that improved awareness and care of psychiatric comorbidities such as substance use disorders may help reduce serious outcomes associated with ADHD.

While previous research has shown that people with ADHD are at increased risk of premature death, the cause-specific risks and the



potential role of psychiatric comorbidities has remained uncertain. Studying national health registries with a sample of 2 675 615 individuals (86 670, 3.2 percent with ADHD diagnosis) born in Sweden between 1983 and 2009, researchers at Karolinska Institutet were able to examine cause-specific risks in ADHD, including suicide and unintentional injuries (e.g. traffic accidents). The researchers also analysed the potential role of early-onset (such as <u>autism spectrum</u> <u>disorders</u>) and late-onset (such as substance use disorders) psychiatric comorbidities.

When analysing children and adults separately, the study revealed that the risk of premature death was higher among adults with ADHD than among children. Individuals who received their ADHD diagnosis later in life also presented with a higher mortality risk compared to those who received their diagnosis earlier. Unintentional injuries and suicide were the leading causes of death among individuals with ADHD, accounting for 35.8 percent and 31.4 percent of the 414 deaths in the ADHD group, respectively.

According to the researchers, the study demonstrates in a variety of ways that psychiatric comorbidity plays an important role in the increased risk of premature death. For example, early-onset psychiatric comorbidity contributed primarily to the association with death due to natural causes such as somatic diseases, whereas later-onset <u>psychiatric comorbidity</u> mainly influenced death due to unnatural causes, including suicide and unintentional injuries.

"Improved understanding of the role of psychiatric comorbidities for associations between ADHD and premature death could facilitate surveillance, intervention and prevention efforts," says Shihua Sun, a doctoral student at the Department of Medical Epidemiology and Biostatistics, Karolinska Institutet and lead author of the study.



Substance use <u>disorders</u> contributed substantially to the risk of death due to suicide, while the increased risk of death due to unintentional injuries seems to be present also in people with only ADHD and no psychiatric comorbidities.

"For individuals with a diagnosis of ADHD and their family members we need to point out that most individuals with ADHD will not suffer from these serious outcomes," says senior author Henrik Larsson, professor at Örebro University and visiting professor at Karolinska Institutet. "The take-home message is that clinicians should consider psychiatric comorbidities carefully in their risk assessments as it may assist in identifying individuals with increased risk of premature <u>death</u>. Increased attention and care of psychiatric comorbidities such as <u>substance use disorders</u> may help reduce serious outcomes associated with ADHD."

More information: Shihua Sun et al. Association of Psychiatric Comorbidity With the Risk of Premature Death Among Children and Adults With Attention-Deficit/Hyperactivity Disorder, *JAMA Psychiatry* (2019). DOI: 10.1001/jamapsychiatry.2019.1944

Provided by Karolinska Institutet

Citation: Psychiatric comorbidity contributes to increased mortality in ADHD (2019, August 8) retrieved 30 April 2024 from https://medicalxpress.com/news/2019-08-psychiatric-comorbidity-contributes-mortality-

adhd.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.