

# The sex gene SRY and Parkinson's disease: How genes act differently in male and female brains

August 14 2019, by Jenny Graves

---



Credit: Monstera Production from Pexels

Parkinson's disease, a debilitating neurodegenerative disease common in elderly people, is [twice as prevalent](#) in men than in women.

A new study [published this month](#) suggests the sex gene (SRY on the male-specific Y chromosome) plays a role in the loss of dopamine-making neurons that underlies this disease.

As well as providing a spectacular example of how genes act differently in male and female brains, this discovery may lead to a new treatment option for men suffering from Parkinson's disease.

## Sex and disease

Many diseases are more common in one sex than the other. For example, multiple sclerosis and other immune disorders are more common in women than men. Parkinson's disease, and several mental health conditions such as schizophrenia and autism, are more common in men than women.

Treatments, too, may be differently effective in men and women because of [differences in expression of genes](#) important for drug metabolism.

The bases of these sex differences are often unclear. Is it a hormonal difference that makes men and women differently susceptible to diseases, and differently amenable to treatment? For instance, the sex difference in Parkinson's disease was previously attributed solely to the protective effect of the hormone oestrogen in female brains.

But as well as hormonal differences, we now have reason to believe genes on sex chromosomes may directly affect the brain.

## Parkinson's disease

Parkinson's disease is a growing problem, particularly with an ageing

population. Nearly [one in 300](#) Australians live with Parkinson's disease. It usually appears in later life as problems in starting and maintaining voluntary movements, and may be accompanied by severe tremor.

Parkinson's disease is caused by a loss of neurons responsible for making dopamine, a hormone and neurotransmitter that sends messages to other nerve cells. Symptoms appear when [70% of these dopamine-synthesising cells](#) have been depleted. We don't understand how these neurons are lost, but expect the effect of loss on motor function is due to the curtailed dopamine production.

Parkinson's disease is progressive and incurable, but the symptoms may be ameliorated and delayed by medications that boost dopamine or substitute for it.

## **SRY and Parkinson's disease**

In humans and other mammals, females have two X chromosomes (XX), and males a single X and a male-specific Y chromosome (XY). SRY is [the master gene](#) on the Y chromosome that determines the male sex of a baby in the embryo.

But research has found SRY seems to be active in other parts of the body, too. In mice and rats, SRY is active in the brain, and in humans it's expressed [in several tissues and organs](#), including the brain.

SRY has been found to be expressed at [abnormally high levels](#) in the brains of mice and rats mutated to have symptoms of Parkinson's disease, and in animals where the disease was induced by chemical treatment.

Previous work showed overactivity of the SRY gene destroys neurons that synthesise dopamine. We're not entirely sure how this happens, but

given the link between dopamine production and Parkinson's disease, it might partly explain why Parkinson's disease affects males more commonly than females.

This [new study](#) now shows that interfering with SRY expression in the brains of rodents with Parkinson's disease ameliorates the severity of symptoms. Vince Harley and Joohyung Lee from the Hudson Institute in Melbourne found that quashing SRY action prevented or mitigated the reduced mobility of male animals with Parkinson's disease.

So, suppressing the activity of SRY in neurons of Parkinson's disease patients could [ameliorate their symptoms](#).

This sort of a cure may be many years away, but it would have a huge impact on the quality of life of thousands of men in Australia living with Parkinson's disease.

## Sex and the brain

Male and female brains really are different at every level; molecular, cellular, and behavioural. For 60 years this has been attributed to sex hormones. But we're beginning to find that genes may also have direct effects.

A recent analysis of the activity of most of the 20,000-odd genes in the bodies of hundreds of men and women showed that more than one-third were expressed much more highly in one sex than the other. This sex bias was not limited to sex organs, but was obvious at many other sites, including the brain.

The effect of SRY in the [brain](#) is a strong demonstration that male and female brains are genetically different in health and disease, and a reminder we must take account of sex differences in diagnosing and

treating [disease](#) in men and women.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: The sex gene SRY and Parkinson's disease: How genes act differently in male and female brains (2019, August 14) retrieved 2 May 2024 from <https://medicalxpress.com/news/2019-08-sex-gene-sry-parkinson-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--