

# Transgender women case study shows sperm production is possible but not certain

August 5 2019

---

Scientists at Magee-Womens Research Institute (MWRI), collaborating with clinicians at UPMC Magee-Womens Hospital and UPMC Children's Hospital of Pittsburgh report two cases in which young transgender women attempted to recover their fertility after starting and stopping gender-affirming medications.

The study, published today in *Pediatrics*, found that one transgender woman was able to produce viable [sperm](#) after a few months of discontinuing her puberty-halting medication, whereas a different patient on hormone therapy could not produce sperm during the time she could psychologically tolerate being off her medication.

"We were interested in examining the timeline for getting viable sperm after stopping masculinity-suppressing medication," said lead author Hanna Valli-Pulaski, Ph.D., a research assistant professor at MWRI.

"Going on and off gender-affirming medications can cause [psychological distress](#) in this population and it's important patients have a discussion with their [health care provider](#) before starting or stopping any treatment."

The research team examined medical records of two [transgender women](#) who tried to preserve their sperm after stopping [hormone therapy](#) and compared their semen quality against eight other transgender women who elected to preserve their sperm before beginning therapy. All of the study participants came through the Fertility Preservation Program in Pittsburgh between 2015 and 2018 as young adults.

One of the patients who elected to preserve their sperm after beginning therapy had been taking the drug Lupron—a sex hormone blocker that halts puberty when taken in adolescence—for six months. She elected to stop taking Lupron to attempt sperm cryopreservation.

Five months later, she was able to produce a sperm sample comparable to those collected from the eight transgender women who saved their sperm prior to undergoing treatment.

Although this one case shows that it's possible to recover sperm after starting gender-affirming therapy, stopping medication for even just a few months can be psychologically distressing, Valli-Pulaski said. For male-to-female transgender individuals, facial hair can start to sprout and the voice begin to deepen after just a few months of stopping [medication](#). It's possible to reverse these effects, but it would take time.

What's more, a second case included in this study showed that fertility doesn't always return quickly after going off gender-affirming drugs.

This patient had been taking estradiol and spironolactone for more than two years. Four months after stopping treatment, she was still unable to produce viable sperm, and at that point, she decided to stop trying for fertility preservation and proceeded with gender reassignment surgery.

The sperm production results of the two study participants provide valuable information that clinicians can share with future patients wishing to have biological children after beginning gender-affirming [therapy](#), notes Valli-Pilaski.

"Right now, there's not much information available about fertility preservation for [transgender](#) patients," Valli-Pulaski said. "If you have any data, it's important to share so that patients, researchers and clinicians can learn from it."

Provided by University of Pittsburgh

Citation: Transgender women case study shows sperm production is possible but not certain (2019, August 5) retrieved 24 April 2024 from <https://medicalxpress.com/news/2019-08-transgender-women-case-sperm-production.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.