

Harms and benefits of estrogen therapy among women with a hysterectomy appear to depend on age when therapy was started

September 10 2019

Patients who started estrogen over the age of 60 did not derive a mortality benefit and experienced more adverse effects particularly among those 70 and older, regardless of whether they had their ovaries removed.

While no statistically significant benefit was seen in the initial study average follow-up period of 7 years, [women](#) in their 50s who had a hysterectomy and their ovaries removed appear to experience a mortality benefit from [estrogen therapy](#) during long-term follow-up (average of 18 years) and did not seem to have harmful health effects. The same was not true for women who started estrogen therapy in their 60s or 70s, whether or not they still had their ovaries. Findings from a randomized [double-blind trial](#) are published in *Annals of Internal Medicine*.

About one third to one half of women who undergo hysterectomy also have their ovaries removed (a procedure called bilateral salpingo-oophorectomy) at the same time to reduce their risk for ovarian cancer in the future. These procedures lead to early menopause and loss of estrogen, which is associated with other health risks. This is the first study to assess, in a randomized trial setting, whether health outcomes of menopausal estrogen therapy differ between women with or without their ovaries.

In a study led by researchers at Brigham and Women's Hospital, Harvard

Medical School, 9,939 women aged 50 to 79 years with prior hysterectomy were included to examine whether estrogen therapy outcomes differed by whether or not the participants' ovaries were also removed. They also assessed whether the effect was different based on the age at which the women began [estrogen therapy](#). Women were randomly assigned to receive estrogen hormone supplements or placebo pills and the researchers documented whether women developed heart disease, breast cancer, or death from any cause, among other measures. They found that menopausal estrogen-alone therapy for a median of 7.2 years was associated with reduced mortality over the long-term (18 years) and had a generally favorable safety profile when initiated before the age of 60 in women who had their ovaries removed. Whether or not their ovaries were intact, [older women](#) did not derive this benefit. Women who had their [ovaries](#) removed and were above age 70 experienced adverse effects of treatment.

More information: Study:

<http://annals.org/aim/article/doi/10.7326/M19-0274>

Patient Summary: <http://annals.org/aim/article/doi/10.7326/P19-0011>

Provided by American College of Physicians

Citation: Harms and benefits of estrogen therapy among women with a hysterectomy appear to depend on age when therapy was started (2019, September 10) retrieved 3 May 2024 from <https://medicalxpress.com/news/2019-09-benefits-estrogen-therapy-women-hysterectomy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.