

One daily combo pill helps lower heart disease risk in study of underserved patients

September 18 2019



Credit: CC0 Public Domain

Taking one daily pill that combined medications to treat high blood

pressure and high cholesterol lowered heart disease risk among underserved patients better than taking several separate medications to treat these risk factors, according to new research published in the *New England Journal of Medicine* and funded by the American Heart Association, the leading voluntary health organization devoted to a world of longer, healthier lives.

Researchers found that patients were also likely to stick to a medication routine if they only had the one pill to take.

"Patients face a variety of barriers to getting the care they need," said study author Daniel Muñoz, M.D, M.P.A., assistant professor of cardiology at Vanderbilt University Medical Center, Nashville, Tennessee. "Those barriers can include cost and complexity of [medication](#) regimens, so innovative strategies are needed to improve the delivery of preventive care, especially when it comes to socio-economically vulnerable individuals."

Muñoz and colleagues studied how a [polypill](#) including four medicines known to lower [high blood pressure](#) and [low-density lipoprotein](#) (LDL) cholesterol would impact patients with high blood pressure and [high cholesterol](#) at a community health center in Mobile, Alabama. Three quarters of the 303 patients studied reported making less than \$15,000 a year; 96% of the participants were black; 60% were women; and the average age was 56.

Half of the patients were given a daily polypill that contained four low-dose medications known to reduce high blood pressure and/or high LDL cholesterol: atorvastatin (10 mg), amlodipine (2.5 mg), losartan (25 mg) and hydrochlorothiazide (12.5 mg). The other half of patients received their usual medical care and medications. Clinicians conducted a medical exam and measured patients' blood pressure and blood cholesterol at the study's start, at two months and at one year.

The researchers found patients taking the polypill had higher reductions in systolic blood pressure and LDL cholesterol compared to those in the usual care group.

In the polypill group, [systolic blood pressure](#) fell an average 9 mm Hg compared 2 mm Hg by one year in the usual care group.

LDL cholesterol fell an average of 15 mg per deciliter in the polypill group versus 4 mg per deciliter in the usual care group.

These differences translate to about a 25% reduction in the risk of experiencing a cardiovascular event, such as a heart attack, according to Muñoz.

Medication adherence over the 12-month study period was high (86%) in patients assigned to the polypill group, compared to past studies that found fewer than half of patients are still on their prescribed blood pressure medicines at one year, according to Muñoz.

"When it comes to preventing cardiovascular disease, simple strategies like the polypill may offer key advantages for patients who face barriers to accessing medical care," Muñoz said. "Simplicity is a big advantage of the polypill. It's once daily; easy to understand; and doesn't require adjustment. Patients are more likely to take their medications as prescribed, which is good for them and their frontline providers."

The population of this study is of particular interest because underserved patients at community health centers tend to be understudied in clinical trials, Muñoz said.

"We need to better understand what works and what doesn't in these settings so we can improve outcomes for people who may be the most vulnerable," he said. "We'll also need to study whether benefits of the

polypill apply to the general population. These are important questions in the conversation about creative, innovative new strategies for the prevention of cardiovascular disease."

The study was undertaken as part of the American Heart Association's Strategically Focused Research Network for Prevention, a \$15 million investment in four research teams —Vanderbilt University Medical Center, Icahn School of Medicine at Mount Sinai, University of Texas Southwestern Medical Center and Northwestern Medical Center. The initiative launched in July 2014 to engage multidisciplinary research teams in finding new innovations for preventing heart disease and stroke.

More information: *New England Journal of Medicine* (2019). [DOI: 10.1056/NEJMoa1815359](https://doi.org/10.1056/NEJMoa1815359)

Provided by American Heart Association

Citation: One daily combo pill helps lower heart disease risk in study of underserved patients (2019, September 18) retrieved 2 May 2024 from <https://medicalxpress.com/news/2019-09-daily-combo-pill-heart-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--