


Dengue virus becoming resistant to vaccines and therapeutics due to mutations in specific protein

September 20 2019



Micrographs of DENV2 with different distribution of smooth and bumpy surfaced particles when incubated at different temperatures to mimic the physiological temperature in the mosquitoes (29°C, left panel), human host (37°C, middle panel) and when humans are experiencing high fever (40°C, right

panel). Colour gradient from blue to red shows the rise in temperature. Credit: Xin-Ni Lim, Emerging Infectious Diseases Programme, Duke-NUS Medical School

Researchers from Duke-NUS Medical School (Duke-NUS), in collaboration with the Agency for Science, Technology and Research (A*STAR)'s Bioinformatics Institute (BII), and the University of Texas Medical Branch (UTMB), U.S., have discovered that the dengue virus changes its shape through mutations in Envelope (E) protein to evade vaccines and therapeutics. The study also gives insights on the types of treatment strategies to use at different stages of infection. This could give rise to new approaches in vaccine development and treatment for dengue disease.

Dengue virus (DENV) infects about 400 million people annually around the world, with a high prevalence in tropical and sub-tropical regions. The virus causes diseases ranging from mild [dengue](#) fever to severe dengue hemorrhagic fever and dengue shock syndrome.

DENV2 exists as smooth spherical surface particles while growing at the mosquito's physiological temperature (29 degrees Celsius). It then changes to bumpy surfaced particles at human physiological temperature (37 degrees Celsius). This ability to morph helps the virus to evade the immune system of the human host. Hence, understanding the mechanism behind this is important for therapeutics and [vaccine](#) development.

"Together with Professor Pei-Yong Shi from UTMB, we found that in laboratory developed DENV2 strains, mutations in the virus' E protein causes its transformation into bumpy particles. These structural changes can cause vaccines and therapeutics to be ineffective against the virus," said Ms Xin-Ni Lim, the study's lead author who is from Duke-NUS'

Emerging Infectious Diseases (EID) Programme.

The team also tested four DENV2 strains obtained from patients. They observed that in contrast to the laboratory adapted viruses, the majority of these clinical strains maintained smooth surface structure at 37 degrees Celsius. However, at 40 degrees Celsius, the temperature of a fever, all virus strains took on a bumpy surface.

"Our study gives a new direction to vaccine development and treatment for dengue disease. For prevention of disease through vaccines that are administered to the patient before dengue infection, we should use those that are effective against the smooth surface virus. When it comes to patients displaying fever symptoms, treatment strategies effective against the bumpy surface particles should be implemented," said Dr. Sheemei Lok, Professor, Duke-NUS' EID and corresponding author of this study.

"This study is a first step towards gaining more insight into how DENV2 reacts and adapts to the host's immunological defenses. We were also able to use computational modeling approaches to predict why particles from different DENV2 strains are more or less adept at morphing from the smooth to bumpy structures. By better understanding the interactions between the [virus](#) and the host, we will be able to develop better therapies and vaccines to treat or prevent infections, and contribute to public health outcomes," said Dr. Peter Bond, Principal Investigator from A*STAR's BII.

The study's findings also show that the lab adapted DENV2 may not be a good model for research, as its structure is different from the clinical strains isolated from patients. The team is planning to study the other DENV serotypes to find out if there are any other possible structural changes.

More information: Xin-Ni Lim et al. Molecular basis of dengue virus

serotype 2 morphological switch from 29°C to 37°C, *PLOS Pathogens* (2019). [DOI: 10.1371/journal.ppat.1007996](https://doi.org/10.1371/journal.ppat.1007996)

Provided by Duke-NUS Medical School

Citation: Dengue virus becoming resistant to vaccines and therapeutics due to mutations in specific protein (2019, September 20) retrieved 19 April 2024 from <https://medicalxpress.com/news/2019-09-dengue-virus-resistant-vaccines-therapeutics.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--