

A single dose of yellow fever vaccine does not offer lasting protection to all children

September 20 2019



Credit: CC0 Public Domain

Yellow fever is a viral infection spread by various species of mosquito and is rife in 34 countries in Africa and 13 in Latin America. Infection may be asymptomatic and go unnoticed or, on the contrary, it may

progress rapidly to severe illness with fever, headache, muscle pain, nausea, vomiting and fatigue. The virus attacks the liver cells, often causing jaundice from which the disease gets its name. Severe bleeding occurs in 25 to 50 percent of cases, with high levels of mortality observed seven to 10 days after the onset of symptoms.

Since 2013, WHO recommends a single dose of the [vaccine](#) for life-long protection. This recommendation is based on proof of long-term efficacy, in vitro and in vivo, established in adults and [children](#) over two years of age. But data on the long-term efficacy of primary vaccination in infants are absent, despite nine- to 12-month-olds being the main targets of routine vaccination in countries in which yellow [fever](#) is endemic. In this respect, WHO recommended research into the long-term persistence of the immunity conferred by vaccination in this age group. This research was performed by José Enrique Mejía from Unit 1043 Center for Pathophysiology of Toulouse Purpan in partnership with Cristina Domingo from Robert Koch Institute in Berlin, and researchers from the U.S., Ghana and Mali.

Their study verified whether children to whom the vaccine was administered at around nine months of age were still protected several years later. The team studied two cohorts, one from Mali (587 children) and the other from Ghana (436 children), in whom the levels of specific antibodies to the [yellow fever](#) virus had been measured four weeks after vaccination. They then repeated the measurement several years later, with findings from previous studies enabling them to estimate that levels above 0.5 IU/ml should protect children from infection.

In the Malian cohort, 4.5 years after vaccination, only half of the children continued to present levels of antibodies above 0.5 IU/ml. And 19.3 percent presented detectable antibodies, but at levels below this recommended threshold (

Citation: A single dose of yellow fever vaccine does not offer lasting protection to all children (2019, September 20) retrieved 3 May 2024 from <https://medicalxpress.com/news/2019-09-dose-yellow-fever-vaccine-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.