

Flavoring ingredient exceeds safety levels in e-cigarettes and smokeless tobacco: study

September 16 2019



Credit: CC0 Public Domain

A potential carcinogen that has been banned as a food additive is present in concerning levels in electronic cigarette liquids and smokeless tobacco products, according to a new study from Duke Health.

The chemical—called pulegone (pronounced pju-leh-goan) - is contained in menthol and mint flavored e-cigarettes and smokeless [tobacco](#) products. Because of its carcinogenic properties, the U.S. Food and Drug Administration banned pulegone as a [food additive](#) last year in response to petitions from [consumer groups](#).

Yet the agency does not regulate the chemical's presence in e-cigarettes and smokeless tobacco, which are promoted as safer alternatives to regular cigarettes.

"Our findings suggest that the FDA should implement measures to mitigate pulegone-related health risks before suggesting mint- and menthol-flavored e-cigarettes and smokeless tobacco products as alternatives for people who use combustible tobacco products," said Sven-Eric Jordt, Ph.D., a professor of the Department of Anesthesiology at Duke and lead author of a study publishing online Sept. 16 in *JAMA Internal Medicine*.

Jordt and research partner Sairam V. Jabba became interested in the topic because the U.S. Centers for Disease Control and Prevention published studies showing that mint- and menthol-flavored [e-cigarette](#) liquids and [smokeless tobacco products](#) marketed in the U.S. contain substantial amounts of pulegone.

The two researchers analyzed whether several top brands of regular menthol cigarettes, three e-cigarette brands, and one smokeless tobacco brand contain enough pulegone to be a cause for concern. They compared the CDC-reported amounts of pulegone with the FDA's exposure risk data—the levels at which exposure-related tumors were reported in animal studies.

Their analysis found that the levels in the e-cigarettes and smokeless tobacco exceeded the thresholds of concern. Regular menthol cigarettes

contained levels below the thresholds.

"Our analysis suggests that users of mint- and menthol-flavored e-cigarettes and smokeless tobacco are exposed to pulegone levels higher than the FDA considers acceptable for intake in food, and higher than in smokers of combustible menthol cigarettes," Jordt said.

"The tobacco industry has long known about the dangers of pulegone and has continuously tried to minimize its levels in menthol cigarette flavorings, so the levels are much lower in [menthol](#) cigarettes than in electronic cigarettes," Jordt said. E-cigarette manufacturers may be less familiar with the dangers and use cheaper ingredients to lower costs.

One limitation of the study is that the FDA's exposure risk calculations are based on oral exposure in animal studies. These risks may apply to the oral exposure through [smokeless tobacco](#) but may differ from inhalation exposure through e-cigarette vapor. There is no toxicity data available on exposure via inhalation. This is concerning because toxicologists consider the lung to be more sensitive to toxic chemicals than the digestive tract.

Provided by Duke University Medical Center

Citation: Flavoring ingredient exceeds safety levels in e-cigarettes and smokeless tobacco: study (2019, September 16) retrieved 5 May 2024 from <https://medicalxpress.com/news/2019-09-flavoring-ingredient-safety-e-cigarettes-smokeless.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--