


New insights on brain connections that are disrupted in patients with coma

September 4 2019



Credit: CC0 Public Domain

Millions of people worldwide suffer from traumatic brain injury each year, and in some severe cases, injured individuals arrive at the hospital in a coma. New research led by investigators at Massachusetts General

Hospital (MGH) sheds light on which connections between brain regions may be severed in such cases. The findings are published in the September 4, 2019, issue of *Neurology*, the medical journal of the American Academy of Neurology.

"We implemented an advanced MRI technique on a clinical MRI scanner to image connections deep within the [brain](#) between areas classically thought to play a role in consciousness, and found evidence of disruption in comatose patients," said lead author Samuel Snider, MD, a clinical fellow in Neurocritical Care at MGH.

Specifically, imaging tests conducted on 16 patients with acute severe [traumatic brain injury](#) who were comatose and 16 matched healthy controls provided an unprecedented three-dimensional view of the brain that revealed a disruption of specific neuronal pathways connecting the brainstem to the thalamus and hypothalamus in comatose patients. It has been known for decades that these pathways are important for consciousness in animals and humans, but no studies have been conducted in the immediate period after brain [injury](#) to examine whether disruption of these pathways is associated with coma.

The findings will be useful because predicting which patients will recover consciousness, and which will remain unconscious or severely disabled, is a major challenge in the intensive care unit. "Clinicians currently lack tools to identify the neural network connections that have been spared from injury and that could support recovery. By mapping these connections with high-resolution MRI while patients are in the [intensive care unit](#), we aim to provide families with a clearer picture of their loved ones' chances of regaining consciousness," said senior author Brian Edlow, MD, director of the Laboratory for NeuroImaging of Coma and Consciousness and associate director of the Center for Neurotechnology and Neurorecovery at MGH.

The researchers noted that all the comatose patients in the study eventually regained consciousness, indicating that they either had a relatively minor injury to this network or reestablished network function with the remaining structural connections. "Sorting out these recovery mechanisms is a focus of our lab, both to provide prognostic guidance for families and therapeutic targets for patients," said Snider.

The investigators also hope to use their technique to provide additional insights on brain biology. "The imaging methods that we implemented in this study provide an opportunity to advance the study of human coma and [consciousness](#)," said Edlow.

More information: *Neurology* (2019). [DOI: 10.1212/WNL.00000000000008163](#)

Provided by Massachusetts General Hospital

Citation: New insights on brain connections that are disrupted in patients with coma (2019, September 4) retrieved 3 May 2024 from <https://medicalxpress.com/news/2019-09-insights-brain-disrupted-patients-coma.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|