

Research shows racial disparities in pregnant women on dialysis

September 26 2019

Pregnancy is not common in women on dialysis due to impaired fertility. New research from the University of Cincinnati finds that among patients with certain kidney disease there is a racial disparity as to who is more likely to become pregnant. Native American, Hispanic and black women have a higher likelihood of pregnancy than white women.

The study, published in the *Journal of the American Society of Nephrology*, finds that compared to [white women](#) on dialysis, Native American [women](#) were 77% more likely, Hispanic women were 51% more likely and [black women](#) were 33% more likely to become pregnant.

"Childbearing is an integral part in women's life," says Silvi Shah, assistant professor in the Division of Nephrology, Kidney CARE (Clinical Advancement, Research and Education) Program at UC and lead author of the study. "However, end-stage [kidney disease](#) or ESKD disrupts this critical element due to impaired fertility, making pregnancy uncommon in women on dialysis. Those who become pregnant face the risk of adverse maternal and fetal outcomes. Since data is so scarce for pregnancy in women undergoing dialysis, we examined the rates, racial differences and factors associated with pregnancy in this high-risk population."

Shah says the study is unique in that it addresses a comprehensive racial group of patients from 2005-13 from a national registry to better understand the incidence of pregnancy and factors associated with it

among dialysis patients. The study further took into account patients with complete Medicaid coverage, thus avoiding the potential shortfalls of registries dependent on voluntary reporting or patient recall. This shows for the first time that pregnancy rates in women with end-stage kidney disease on dialysis are higher than in previous reports.

"This is telling us that there are additional factors, which could be biological, contributing to these racial differences. While we can speculate that [health literacy](#) has an important role, the real reasons remain unknown," says Shah. "However, this information will help patients and [health care providers](#) in shared decision-making regarding management of their reproductive health."

Assisting in the research were Annette Christianson and Karthikeyan Meganathan, research associates in the UC Department of Biomedical Informatics; Anthony Leonard, research associate professor in the UC Department of Family and Community Medicine; Daniel Schauer, of the UC Department of Internal Medicine and Charuhas Thakar, director, Division of Nephrology Kidney CARE Program. Shah is supported by the intramural funds from the UC Division of Nephrology.

The study evaluated 47,555 women of childbearing age who were undergoing dialysis between Jan. 1, 2005, and Dec. 31, 2013, using data from the United States Renal Data System. Overall, 2,352 pregnancies were identified. The mean age of the study population at transplant was approximately 33 years. The rate of pregnancy in dialysis patients was 17.8 per thousand person-years (PTPY) and highest rate was in women aged 20-24 (40.9 PTPY).

Pregnancy rates were highest in Native American women followed by Hispanics, blacks, Asians and whites. Patients with diabetes as the cause of ESKD had the lowest pregnancy rates and [pregnancy](#) was more likely in women with ESKD due to malignancy, glomerulonephritis and

hypertension.

More information: Silvi Shah et al, Racial Differences and Factors Associated with Pregnancy in End Stage Kidney Disease Patients on Dialysis in the United States, *Journal of the American Society of Nephrology* (2019). [DOI: 10.1681/ASN.2019030234](https://doi.org/10.1681/ASN.2019030234)

Provided by University of Cincinnati

Citation: Research shows racial disparities in pregnant women on dialysis (2019, September 26) retrieved 7 May 2024 from <https://medicalxpress.com/news/2019-09-racial-disparities-pregnant-women-dialysis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.