


Scientists discover how cancer cells regenerate in the prostate

September 16 2019



Treating patients with an inhibitor of Notch in addition to radiotherapy would improve prostate cancer treatment, the researchers say. Credit: University of York

Scientists have discovered a mechanism by which cancer cells in the prostate can regenerate following radiotherapy—paving the way for improved prostate cancer treatment.



The study, carried out by researchers and the University of York and funded by York Against Cancer, identified signals which can prompt cancer cells to grow again immediately after plasma therapy or radiotherapy—causing the treatments to fail.

The scientists, from the Department of Biology and the York Plasma Institute, discovered that while most <u>cancer cells</u> were likely to be killed off by treatment, cell signals within a process known as the "Notch response' can kick-start growth in a small number of cancer stem cells, resulting in regeneration of both cancer and normal tissues.

Triggered

The team designed experiments to observe <u>prostate cancer</u> cells every two hours, discovering that these signals were triggered between four and eight hours after treatment. By 72 hours after treatment, the targeted cells were either dead or growing again.

Treating patients with an inhibitor of Notch in addition to radiotherapy would prevent cancer cells from re-growing, the researchers say.

Prostate cancer is the most common male cancer in the UK—one in eight men will get it in their lifetime.

Preventing relapse

Professor Norman Maitland, from the Department of Biology at the University of York, said: "The strong response we observed was due to the powerful cellular function called Notch. It was present in both normal and cancer cells and it's how a cell population can regenerate itself after radiotherapy.



"Radiotherapy and, potentially, Low Temperature Plasma are really effective ways to kill cancer cells. The problem always is that a proportion of the cancers relapse. By unpicking the responses of cells from men with prostate cancer in Yorkshire we have unearthed a solution.

"The implication is that if you were able to treat cells with both an inhibitor of Notch and radiotherapy you would stop the regenerating response. That's precisely what we were able to do—it is an important piece of research and that one that I am particularly proud to have been involved in during my 28-year career in York."

Common cancers

The research may pave the way for <u>clinical trials</u>, which could take place more quickly than usual because the drugs used in the lab tests are chemotherapy agents already approved for use in humans.

Professor Maitland said that while the research had concentrated on prostate cancer cells, it could also have ramifications for other common cancers such as <u>breast cancer</u>, although a slightly different response had resulted from research on breast cancer cells.

More information: J.R. Packer et al. Notch signalling is a potential resistance mechanism of progenitor cells within patient-derived prostate cultures following ROS -inducing treatments, *FEBS Letters* (2019). <u>DOI:</u> 10.1002/1873-3468.13589

Provided by University of York



Citation: Scientists discover how cancer cells regenerate in the prostate (2019, September 16) retrieved 19 April 2024 from

https://medicalxpress.com/news/2019-09-scientists-cancer-cells-regenerate-prostate.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.