

Shoulder pain surgery: One popular procedure not backed by evidence

September 13 2019, by Tim Jones and William Hollingworth



Credit: AI-generated image (disclaimer)

Many countries have strict regulations to ensure new drugs are effective and are worth the money before doctors can prescribe them. But surgical procedures are often <u>less strictly regulated</u>. This can lead to people having risky surgery even though there is no clear evidence that it works. One such surgery is called subacromial decompression—one of the most



common procedures to treat chronic shoulder pain.

Up to <u>two-thirds</u> of people experience shoulder pain during their life. This is often caused by bony spurs forming on the shoulder blade, which can lead to swelling in the surrounding area. Subacromial <u>decompression</u> is keyhole surgery to remove these spurs.

Several randomized control trials—the gold standard for <u>clinical</u> <u>evidence</u>—have found no evidence that subacromial decompression is effective. Two recent high-quality randomized controlled trials (RCTs) in the UK (<u>CSAW</u>) and Finland (<u>FIMPACT</u>) found that subacromial decompression is no more effective than placebo surgery—where the surgeon makes a small cut but doesn't remove the spurs.

And a <u>recent review</u> of the evidence, which included the CSAW and FIMPACT studies, found high-quality evidence that subacromial decompression doesn't improve pain, movement or health-related quality of life. Following this, a <u>BMJ article</u> strongly recommended that surgeons stop performing this surgery for chronic shoulder pain.

<u>We recently</u> investigated the extent to which this surgery is performed, given the lack of evidence to support its use. For our study, we examined the relationship between research evidence and treatment. We found that <u>health systems</u> around the world have been carrying out a lot of subacromial decompression surgery over the past 25 years.





Credit: AI-generated image (disclaimer)

Over ten years, the number of subacromial decompression procedures nearly doubled in England, from 15,112 procedures (30 per 100,000 people) in 2007-08 to 28,802 procedures (52 per 100,000 people) in 2016-17. Surgery rates were generally more than twice as high (102 to 131 per 100,000 people) in other regions (New York State, Western Australia, Florida State and Finland).

The example of subacromial decompression highlights that, without timely high-quality research, the use of surgery that doesn't work can increase over time. It is important to identify these procedures early and run trials to examine their value. Healthcare systems need effective methods to stop surgery that doesn't work and isn't value for money.



Not the only surgery that doesn't work

The situation with subacromial decompression surgery is not unique. Another example is the <u>use of stents</u> to open narrowed arteries to treat chest pain (angina). Around half a million people receive stents for chest pain each year in the US and Europe, yet a recent <u>placebo-controlled</u> <u>RCT</u> found no difference in angina <u>pain</u> outcomes between inserting a stent and using standard drugs.

Also, consider keyhole surgery to clean out the knee joint, on which around <u>US\$4 billion</u> is spent each year in the US. Recent RCTs found <u>no</u> <u>evidence</u> of effectiveness to justify the spending.

Given <u>limited resources</u> and the potential risk to patients, doctors should only recommend surgical procedures backed by evidence. We need more robust trials to evaluate <u>surgery</u> where there is currently not enough evidence. A balance needs to be struck between supporting innovation and preventing unnecessary treatment. Health services must develop processes that support this balance, to reduce harm to patients and get the maximum health benefit for our money.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Shoulder pain surgery: One popular procedure not backed by evidence (2019, September 13) retrieved 20 May 2024 from <u>https://medicalxpress.com/news/2019-09-shoulder-pain-surgery-popular-procedure.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.