


# Source water key to bacterial water safety in remote Northern Australia

September 5 2019

---



Credit: CC0 Public Domain

In the wet-dry tropics of Australia, drinking water in remote communities is often sourced from groundwater bores. The geochemistry of that groundwater impacts the occurrence of opportunistic pathogens in the drinking water supply, researchers now report in *PLOS Neglected Tropical Diseases*.

Many aquifers in the tropics of Northern Australia are shallow and contain naturally high levels of [iron](#), which encourages environmental iron bacteria to form biofilms in pipes and the growth of opportunistic pathogens. Some of these bacteria include the emerging threat *Burkholderia pseudomallei* which causes the fatal disease melioidosis in humans and animals. Researchers have not previously studied the microbial composition of water from remote bores.

In the new work, Mirjam Kaestli of Charles Darwin University, Australia and colleagues including Menzies School of Health Research sampled water and biofilms from three remote Indigenous communities in Australia's Northern Territory. All three communities had previously reported *B. pseudomallei* clinical infections. The aquifers in each community varied by their age and depth; in addition, one had low levels of iron, another had average levels of iron, and a third had high levels of iron. Water was collected from five points in each [water system](#), including both untreated sections of the system, and points after chlorination.

On average 19 times more bacterial DNA was found in the source water from the medium iron water supply and 83 times more in the high iron water supply compared to the low iron system. In addition, biofilms were more common in these supplies and bacteria known to cycle iron were found in the medium and high iron water systems. *B. pseudomallei* were also found in these systems and the genomes revealed five different genetic variants of the bacteria, including three that had been previously reported in Australia. Overall, chlorination successfully contained both *B. pseudomallei* and another opportunistic pathogen *Pseudomonas aeruginosa*.

"Results from this study will inform future studies to ultimately improve management guidelines for [water](#) supplies in the wet-dry tropics," the researchers say.

**More information:** Kaestli M, O'Donnell M, Rose A, Webb JR, Mayo M, et al. (2019) Opportunistic pathogens and large microbial diversity detected in source-to-distribution drinking water of three remote communities in Northern Australia. *PLOS Neglected Tropical Diseases* 13(9): e0007672. [doi.org/10.1371/journal.pntd.0007672](https://doi.org/10.1371/journal.pntd.0007672)

Provided by Public Library of Science

Citation: Source water key to bacterial water safety in remote Northern Australia (2019, September 5) retrieved 19 April 2024 from <https://medicalxpress.com/news/2019-09-source-key-bacterial-safety-remote.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.