


## Attachment theory: Type of fixation pins matters during stereotactic radiosurgery

September 3 2019, by Cindy Starr, Msj

---



A stereotactic frame is secured to the head with pins. Credit: Mayfield Clinic

In the field of stereotactic radiosurgery of the brain, the fixation pins that keep a head frame perfectly stable during treatment perform differently.

That was the surprising discovery made by Ronald Warnick, MD, a neurosurgeon with Mayfield Brain & Spine and Co-Director of the Gamma Knife Center at The Jewish Hospital-Mercy Health. Dr. Warnick's [findings](#) were published this month in the *Journal of Radiosurgery and SBRT*.

Stereotactic radiosurgery involves the delivery of high-energy rays to treat tumors and other diseases. Radiation damages DNA within cells, making them unable to divide and grow. When patients with brain lesions are treated with a single, high dose of radiation, their head is immobilized in a stereotactic head frame, which is attached with four identical pins made of either aluminum or titanium.

Dr. Warnick's retrospective study aimed to discover whether there was a difference in penetration of the two types of pins, which the Gamma Knife manufacturer, Elekta, provided. Elekta offered no instructions other than to say that the aluminum pins produced less artifact on CT scans. Dr. Warnick assessed the pins' penetration on 101 adult patients treated with single-dose radiosurgery from July 2013 to June 2018. All of the pins were tightened with the same type of torque wrench and to the same degree. Measurements were made at each of the four pin sites on CT images, and the depth of penetration was graded.

"We had utilized both types of pins, not knowing that there was any difference," Dr. Warnick says.

As it turned out, there was a difference. "I noted that the aluminum pins were not penetrating the outer layer of the skull," Dr. Warnick says. "They were essentially acting as a vice grip. The titanium pins, by comparison, penetrated the outer layer of the skull, which should result in higher frame stability."



Ronald Warnick, MD, neurosurgeon, Mayfield Brain & Spine, and Co-Director of the Gamma Knife Center at The Jewish Hospital-Mercy Health Credit: The Jewish Hospital - Mercy Health

Although there is no clinical evidence that the head frame was capable of movement while in the "vice grip" with the aluminum pins, stability of the frame is recognized as critical during Gamma Knife radiosurgery, when 192 low-dose beams intersect at the tumor or lesion. Gamma Knife technology promises accuracy to .5 millimeters while delivering a radiation dose to healthy tissue that is two to five times lower than competing technologies.

"It seems like an old concept, but there has been a resurgence of interest in the stability of the frame," Dr. Warnick says. "How do we achieve a

stable frame? How do we verify that the frame is stable? After looking at 101 patients—using 404 pins—and finding major differences, we concluded that aluminum pins should only be used in rare circumstances."



Radiation is delivered through 192 openings in the Gamma Knife machine.  
Credit: Mayfield Clinic

Aluminum pins are indicated, Dr. Warnick says, in cases where minimal penetration is desired. Those situations include the presence of 1) a large frontal sinus, located above the ridges of the brow; 2) a prior craniotomy, an opening of the skull that allowed for the removal of a tumor; and 3) a very thin skull.

Dr. Warnick predicts that the study will reverberate because physicians

have been conditioned to think that all pins are appropriate for all patients. "We now know that pins should be selected based on the individual patient," he says. "The end result will be heightened submillimetric accuracy for an already highly accurate Gamma Knife system."

Co-author of the study was Eunsun Yook, MS, of the TriHealth Hatton Research Institute.

**More information:** [www.oldcitypublishing.com/jour...-sbrt-6-3-p-241-246/](http://www.oldcitypublishing.com/jour...-sbrt-6-3-p-241-246/)

Dutta SW, Kowalchuk RO, Trifiletti DM, Peach MS, Sheehan JP, Lerner JM, Schlesinger D. Stereotactic shifts during frame-based image-guided stereotactic radiosurgery: Clinical measurements. *Int J Radiat Oncol Biol Phys* 2018;102(4):895-902.

Toyota S, Seta H, Kubo H, Muramatsu M, Takeda K. Relationships between head fixation pins for radiosurgery and the skull bone: Usefulness of a torque wrench. *Radiat Med* 2003;Mar-Apr;21(2):94-98.

Rojas-Villabona A, Miszkiel K, Kitchen N, Jäger R, Paddick I. Evaluation of the stability of the stereotactic Leksell Frame G in Gamma Knife radiosurgery. *J Appl Clin Med Phys* 2016 May 8;17(3):75-89.

Safae M, Burke J, McDermott MW. Techniques for the application of stereotactic head frames based on a 25-year experience. *Cureus* 2016 Mar 25;8(3):e543.

Provided by Mayfield Clinic

Citation: Attachment theory: Type of fixation pins matters during stereotactic radiosurgery (2019, September 3) retrieved 18 April 2024 from <https://medicalxpress.com/news/2019-09-theory-fixation-pins-stereotactic-radiosurgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.