

# Vigorous exercise once or twice a week linked to best outcomes in heart patients

September 27 2019

---



Credit: CC0 Public Domain

Vigorous physical activity one or two times a week is associated with the lowest death rates in patients with stable coronary artery disease, reports a study published today in the *European Journal of Preventive*

*Cardiology*, a journal of the European Society of Cardiology (ESC).

The findings come ahead of World Heart Day on 29 September, which highlights that [cardiovascular disease](#) is the world's number one killer, causing 17.9 million deaths a year.

Study author Dr. Simone Biscaglia of the University of Ferrara, Italy said: "The optimum level of [exercise](#) is achievable in almost all patients with stable coronary artery disease worldwide—but was only reached in one-third of participants in our study. Walking once or twice a week at a pace that causes shortness of breath or a raised [heart rate](#) is all that's needed."

It is [common knowledge](#) that exercise is healthy for heart patients—but how often and at what intensity is controversial. This analysis of the CLARIFY registry examined these questions. In addition, since most [heart patients](#) do little exercise, the study examined the most important determinants of [physical activity](#).

CLARIFY enrolled 32,370 outpatients with stable coronary artery disease from 45 countries in Africa, Asia, Australia, Europe, the Middle East, and North, Central, and South America who were followed up for five years. Patients were divided into four groups according to self-reported physical activity at the inclusion visit:

1. sedentary (16.1 percent)
2. only light physical activity most weeks (51.4 percent)
3. vigorous physical activity once or twice per week (16.8 percent)
4. [vigorous physical activity](#) three or more times per week (15.7 percent)

The primary outcome was the composite of cardiovascular death, myocardial infarction, and stroke. Compared to the light activity group,

patients who did vigorous exercise one or two times per week had the lowest risk of the primary outcome, with a hazard ratio (HR) of 0.82. More frequent activity did not lead to further benefit. Sedentary patients had the highest risk of the primary outcome (HR 1.32).

Patients in the group performing vigorous exercise one or two times per week also had the lowest risks of all-cause death, cardiovascular death, and stroke—19 percent, 21 percent, and 26 percent lower, respectively, compared to the light activity group.

"Stable coronary artery disease patients should avoid being sedentary," said Dr. Biscaglia. "The goal should be to exercise every week, not to achieve the highest possible frequency, since this is unlikely to be sustainable and does not translate into better outcomes."

Women were less likely to be physically active than men. Dr. Biscaglia said: "In previous studies we found that women were less often referred to cardiac rehabilitation programs and had less intensive lifestyle counseling. Clinicians should provide the same exercise advice and referrals to women as they do for men, which should result in equal levels of physical activity."

Other predictors of low physical activity were comorbidities such as peripheral artery disease, diabetes, and previous myocardial infarction or stroke. "Patients with these conditions may worry that exercise could worsen their symptoms," said Dr. Biscaglia. "But our study shows that they get the same benefit as others with stable coronary artery disease so these conditions should not deter them from being physically active."

The rate of myocardial infarction during follow-up was comparable between the four physical activity groups. "This can reassure patients with stable coronary artery disease—fear of having a heart attack should not be a barrier to physical activity," said Dr. Biscaglia. "More research

is needed to discover what drives the reduction in cardiovascular death seen with exercise—partly this is due to fewer fatal strokes, but the other mechanisms are currently unknown."

**More information:** Simone Biscaglia et al. Relationship between physical activity and long-term outcomes in patients with stable coronary artery disease, *European Journal of Preventive Cardiology* (2019). [DOI: 10.1177/2047487319871217](https://doi.org/10.1177/2047487319871217)

Provided by European Society of Cardiology

Citation: Vigorous exercise once or twice a week linked to best outcomes in heart patients (2019, September 27) retrieved 2 May 2024 from <https://medicalxpress.com/news/2019-09-vigorous-week-linked-outcomes-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--