

Walking speed may predict return to work in young stroke survivors

September 26 2019



Credit: CC0 Public Domain

A simple walking speed test may help predict whether young adult stroke survivors are ready to return to work, according to new research

published in *Stroke*, a journal of the American Stroke Association, a division of the American Heart Association.

Today, one of every four people who have a stroke is under the age of 65, and up to 44% of those may not be able to return to work, mostly due to challenges they experience with walking.

"Stroke affects motor control and motor function. To return to work, you must be able to walk to your car, bus, office and meeting rooms. If you can't walk or you get tired easily, your ability to do your job is going to be seriously impacted," said Hannah Jarvis, Ph.D., lead study author and research associate at Manchester Metropolitan University in Manchester, United Kingdom.

Researchers analyzed the relationship between walking [speed](#) and readiness to return to work after stroke. They compared mobility in 46 [stroke survivors](#) from Wales, U.K. - between the ages of 18 and 65—to 15 people who had not had a stroke. Using a walking test, researchers measured how fast and far participants walked in three minutes. A benchmark measure was calculated that predicted readiness to return to work. They found:

A walking speed of more than 3 feet per second was a critical threshold. Stroke survivors walking slower than this were significantly less likely to return to work.

Of the 23% who returned to work, 90% walked faster than the critical threshold. Those who returned to work walked almost 6 feet per second, compared to those who didn't go back to work and only covered about two and a half feet per second.

Young stroke survivors were less efficient at walking (known as metabolic cost of walking) than their healthy counterparts. This means

that walking at a similar pace requires more effort and results in becoming tired more easily.

"There was a huge range in outcomes in our stroke population, highlighting the fact that some people recover quickly and can return to work, while others are so severely affected that they cannot go back to work," Jarvis said.

"Walking speed is a really useful tool for clinicians to use to predict return to work. It's simple, low cost and effective. In addition, clinicians can use this measure to guide their patients during rehabilitation. For example, they can focus on increasing walking speed and maintaining quality of walking in order to give their patients a chance of going back to work," Jarvis added.

According to the American Stroke Association, someone in the U.S. has a [stroke](#) every 40 seconds on average, and 5.2 million first strokes (31%) are in those younger than 65.

More information: *Stroke* (2019). [DOI: 10.1161/STROKEAHA.119.025614](#)

Provided by American Heart Association

Citation: Walking speed may predict return to work in young stroke survivors (2019, September 26) retrieved 25 April 2024 from <https://medicalxpress.com/news/2019-09-young-survivors.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--