

Youngest in classroom diagnosed more often with ADHD, other problems

September 23 2019, by Serena Gordon, Healthday Reporter



(HealthDay)—If a child can't sit still or blurts out random thoughts in

kindergarten or first grade, does the child have attention-deficit/hyperactivity disorder (ADHD)? Or is the youngster just not mature enough to sit still yet?

Both are possibilities, and whichever might be true, new research suggests that the youngest kids in class are being diagnosed with ADHD, [intellectual disability](#) and even depression around 30% more often than their oldest classmates.

"Children in the same [school] year as each other can be nearly a whole year apart in terms of age. We have known for some time that [children](#) who are young in their year at school tend to do less well academically, and are more likely to be diagnosed with hyperactivity disorders," said study co-author Jeremy Brown, from the London School of Hygiene and Tropical Medicine, in England.

Brown noted that this study wasn't designed to determine the cause for these issues, just to find an association. But he said there are a number of potential reasons for the link.

"Younger children may find it harder to concentrate in class, leading to increased diagnoses of hyperactivity. Inferior academic performance and poorer peer relationships could have negative consequences on mental health," he explained.

For the study, the researchers reviewed electronic health care records from more than 1 million [school children](#) in the United Kingdom.

The investigators found that children born in the last quarter of a school year were 36% more likely than kids born in the first quarter of a school year to be diagnosed with ADHD. The youngest kids were 30% more likely to have an intellectual disability and 31% more likely to have depression than their older classmates.

Children born in the second quarter appeared to have a slight disadvantage compared to the older kids in terms of intellectual disability. And, kids born in the third quarter of the year were 20% more likely than the oldest kids to have an intellectual disability, the findings showed.

Brown said about 800,000 kids start primary school in the United Kingdom each year. "Based on our findings, over the course of their schooling up to age 16 years, we would expect around 500 more of the youngest quarter in the year might be diagnosed with depression, compared to the oldest," he said.

For ADHD, that number was 1,200 more of the youngest children, and for intellectual disability, the difference would be about 500 more of the youngest children with a problem, he added.

Brown said the solution isn't clear, and more research is needed. More support for younger children might be of value, he suggested. It also might help to have a fairer and clearer process for delaying school entry to young and developmentally immature children.

Dr. Meg Benningfield, director of the division of child and adolescent psychiatry at Vanderbilt University in Nashville, Tenn., said, "If you compare a 4-year-old to a 5-year-old, there's a big difference developmentally. When [younger children](#) are held to the same standard as older kids, there's a risk for both overdiagnosis and underdiagnosis." (Just as an over-fidgety kid might be assumed to have ADHD when they don't, the reverse is also true, and teachers might excuse poor behavior in a child because they're young, when they actually have ADHD.)

Benningfield said that if parents are concerned about their child's behavior—no matter their age—they should talk with their child's pediatrician or a child and adolescent psychiatrist, if available.

She recommended helping children who are younger than their class peers focus on how they, themselves, are improving. What can you do today that you couldn't do yesterday? How did you do on your spelling test today compared to last week? Teach them to look at how they're growing and getting better, and minimize comparisons to other children.

Dr. Andrew Adesman is chief of developmental and behavioral pediatrics at Cohen Children's Medical Center in New Hyde Park, N.Y.

He said that previous studies have shown the connection to ADHD and younger age in a grade. But he was surprised by the findings on intellectual disability. Adesman said that IQ test scoring is adjusted based on a child's exact age, so he wouldn't have expected that difference.

The "findings suggest that more consideration should be given to a child's age relative to his or her classmates when concerns about short attention span, impulsivity or restlessness are first raised."

Adesman also said that a child's birthdate shouldn't be the sole factor in deciding when kids start [school](#). He said that a [child](#)'s maturity level should also be considered.

The study was published online Sept. 23 in *JAMA Pediatrics*.

More information: Learn more about an early diagnosis of ADHD from the [Child Mind Institute](#).

Copyright © 2019 [HealthDay](#). All rights reserved.

Citation: Youngest in classroom diagnosed more often with ADHD, other problems (2019, September 23) retrieved 9 April 2024 from <https://medicalxpress.com/news/2019-09-youngest->

classroom-adhd-problems.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.