

In Baltimore, lower income neighborhoods have bigger mosquitoes

October 16 2019



Paul Leisnham, a co-author on the study from University of Maryland - College Park, collecting mosquito larvae in a vacant lot in Baltimore. Credit: Shannon LaDeau.

Low-income urban neighborhoods not only have more mosquitoes, but they are larger-bodied, indicating that they could be more efficient at transmitting diseases. So reports a Cary Institute of Ecosystem Studies-



led study, published in the *Journal of Medical Entomology*, investigating how socioeconomics influences mosquito-borne disease risk in Baltimore, Maryland.

As part of the Baltimore Ecosystem Study, Cary Institute scientists have been investigating how environmental and social conditions regulate mosquito numbers. Tiger mosquitoes (Aedes albopictus) were the focus of this study. This <u>invasive species</u> dominates in urban areas, where it is an aggressive day-biter that targets people and can transmit an array of viruses including <u>dengue fever</u>, Zika, chikungunya, Eastern Equine Encephalitis, and West Nile virus.

Cary Institute disease ecologist Shannon LaDeau, senior author on the study, explains, "More people are living in cities. At the same time, many other species are also adapting to city living. Invasive species like the tiger mosquito increasingly thrive in temperate urban areas, living among us and fundamentally altering the risk of local disease emergence. "

Previous studies have found that wing length is an accurate proxy for body size in mosquitoes, and body size influences traits that are important to disease transmission. LaDeau and colleagues measured wing length of tiger mosquitoes trapped across a continuum of neighborhoods in Baltimore, Maryland. Results where clear: mosquitoes from less affluent blocks were larger than mosquitoes from more affluent blocks.





Average wing length in high-abandonment blocks was larger than lowabandonment blocks. Credit: Grace Katz

Block-to-block differences in <u>abandoned buildings</u>, vacant lots, and trash removal influence mosquito populations, biting patterns, and disease risk. LaDeau explains, "People living in neighborhoods with abandoned infrastructure are more at risk, because tiger mosquitoes flourish in less managed landscapes. They can breed in the water filling a bottle cap or crushed can. Abundant juvenile habitat sets the stage for healthy adult mosquitoes."

Adult female tiger mosquitoes were collected from 13 residential blocks in Baltimore over the course of three summers. Trapping took place in June and July, 2015-2017. Blocks were distributed among five



neighborhoods ranging in socioeconomic status—with varying levels of management and occupancy.

For each block, the team measured the percentage of abandoned structures and counted discarded container habitats—defined as any object that could potentially hold water—to determine whether differences in these features across blocks influenced numbers of juvenile tiger mosquitoes and adult female body size.

Over the three years of the study, 1097 mosquitoes were collected and measured. The team found that mosquitoes from blocks with higher abandonment had larger wings than those collected on more affluent blocks with less abandonment. Wing length differences of less than one millimeter can affect traits like fecundity, longevity, and ability to spread disease.





Tiger mosquitoes (*Aedes albopictus*) lay drought-resistant eggs that remain viable for months. They can breed in very small containers of water. Credit: CDC

They also counted juvenile tiger mosquitos and related these numbers to



the abundance of unmanaged container habitats in a block. Less affluent blocks averaged 400 habitats per square kilometer with 80% containing mosquito larvae. More affluent blocks averaged only 50 containers per square kilometer with less than 5% containing mosquito larvae.

Lead author Grace Katz worked on the project as a high school intern and was responsible for measuring hundreds of wings. She explains, "Understanding how landscape features influence mosquito size and their ability to make us sick is important if we want to better manage mosquitoes in cities."

LaDeau concludes, "The trends we're seeing in mosquito numbers and body size map onto socioeconomic patterns at the block-level. There are environmental justice and equity implications at play. Residents of less affluent neighborhoods are exposed to more mosquitoes. If those mosquitoes have greater longevity or fitness, as previous studies on body size suggest, then residents may be at a greater risk of contracting mosquito-borne illnesses. There's a pressing need to address infrastructure abandonment and waste management to protect all residents."

More information: Grace Katz et al. Aedes albopictus Body Size Differs Across Neighborhoods With Varying Infrastructural Abandonment, *Journal of Medical Entomology* (2019). <u>DOI:</u> <u>10.1093/jme/tjz170</u>

Provided by Cary Institute of Ecosystem Studies

Citation: In Baltimore, lower income neighborhoods have bigger mosquitoes (2019, October 16) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2019-10-baltimore-income-neighborhoods-bigger-</u>



mosquitoes.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.