

New biomarkers for childhood asthma may facilitate early diagnosis

October 21 2019



Credit: CC0 Public Domain

Researchers at Karolinska Institutet in Sweden have followed children who have sought emergency care for wheezing during their first years of life and found specific biomarkers that can predict the need for asthma medication several years later. The study is published in the prestigious *European Respiratory Journal*.

Today, biomarkers for asthma are lacking in the [youngest children](#), although asthma and asthma-like respiratory disorders are one of the most common causes of emergency visits to the hospital at preschool age. Therefore, it is difficult to assess the prognosis and need for treatment for these children.

"Our goal is to develop a molecular tool that can complement today's clinically based diagnosis and prognosis," says Cilla Söderhäll, associate professor of medical genetics at the Department of Women's and Children's Health, Karolinska Institutet, who led the study.

For a number of years, the researchers have followed a group of children who sought care for acute wheezing at the pediatric emergency unit of the Astrid Lindgren Children's Hospital in Stockholm, Sweden. At the time of the emergency visit, the children were between six months and three years of age. In connection with the ongoing wheezing, the researchers were able to identify specific biomarkers that predict lung function and the need for [asthma medication](#) at seven years of age. The biomarkers are based on differences in [gene expression profiles](#) in this patient group compared to children who have not sought [emergency care](#) for wheezing.

In addition, the researchers found an association between Vitamin D levels in infants and [gene expression](#) in acute wheezing, which supports the impact of Vitamin D levels for wheeze.

"Our findings contribute to increased knowledge about asthma in

preschool children and provide information that increases the opportunities to identify children at risk of chronic asthma already at an early emergency visit," says Cilla Söderhäll. "In the long term, early identification of [preschool children](#) at risk of developing chronic asthma can contribute to a more personalized and effective treatment of asthma in [children](#)."

The research was supported by The Swedish Heart-Lung Foundation, Stiftelsen Sven Jerrings fond, the Centre for Allergy Research at Karolinska Institutet, the Swedish Association for Allergology (SFFA), Freemason Child House Foundation in Stockholm, Konsul Th C Berghs stiftelse, the Swedish Asthma and Allergy Association's Research Foundation, EU's seventh framework program, the Swedish Research Council, Knut and Alice Wallenberg Foundation, and Sigrid Jusélius Foundation. Two of the co-authors have a patent pending for "GlobinLock," a method used in the study.

More information: Shintaro Katayama et al. Acute wheeze-specific gene module shows correlation with vitamin D and asthma medication, *European Respiratory Journal* (2019). [DOI: 10.1183/13993003.01330-2019](#)

Provided by Karolinska Institutet

Citation: New biomarkers for childhood asthma may facilitate early diagnosis (2019, October 21) retrieved 25 April 2024 from <https://medicalxpress.com/news/2019-10-biomarkers-childhood-asthma-early-diagnosis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--