

Brain networks more stable in individuals with higher cognitive abilities

October 15 2019



Credit: CC0 Public Domain

The interconnections and communication between different regions of the human brain influence behavior in many ways. This is also true for individual differences in higher cognitive abilities. The brains of more

intelligent individuals are characterized by temporally more stable interactions in neural networks. This is the result of a recent study conducted by Dr. Kirsten Hilger and Professor Christian Fiebach from the Department of Psychology and Brain Imaging Center of Goethe University Frankfurt in collaboration with Dr. Makoto Fukushima and Professor Olaf Sporns from Indiana University Bloomington, U.S. The study was published online in the scientific journal *Human Brain Mapping* on 6th October.

Intelligence and its neurobiological basis

Various theories have been proposed to explain the differences in individuals' cognitive abilities, including neurobiological models. For instance, it has been proposed that more intelligent individuals make stronger use of certain brain areas, that their brains generally operate more efficiently, or that certain brain systems are better wired in smarter people. Only recently have methodological advances made it possible to investigate the temporal dynamics of human brain networks using functional magnetic resonance imaging (fMRI). An international team of researchers from Goethe University and Indiana University Bloomington analyzed fMRI scans of 281 participants to investigate how dynamic [network](#) characteristics of the human brain relate to general intelligence.

Stability of brain networks as general advantage

The human brain has a modular organization—it can be subdivided into separate networks that serve different functions such as vision, hearing, or the control of voluntary behavior. In their current study, Kirsten Hilger and colleagues investigated whether this modular organization of the human brain changes over time, and whether or not these changes relate to [individual differences](#) in the scores that study participants achieved in an intelligence test.

The results of the study show that the modular brain network organization of more intelligent persons exhibits fewer fluctuations during the fMRI measurement session. This increased stability of brain network organization was primarily found in brain systems that are important for the control of attention.

Attention plays a key role

"The study of the temporal dynamics of human brain networks using fMRI is a relatively new field of research" says Hilger. "The temporally more stable network organization in more intelligent individuals could be a protective mechanism of the brain against falling into maladaptive network states in which major networks disconnect and communication may be hampered."

She also stresses that it remains an open question how these network properties influence cognitive ability: "At present, we do not know whether the temporally more stable [brain](#) connections are a source or a consequence of higher intelligence. However, our results suggest that processes of controlled attention—that is, the ability to stay focused and to concentrate on a task—may play an important role for general intelligence."

More information: Kirsten Hilger et al. Temporal stability of functional brain modules associated with human intelligence, *Human Brain Mapping* (2019). [DOI: 10.1002/hbm.24807](https://doi.org/10.1002/hbm.24807)

Provided by Goethe University Frankfurt am Main

Citation: Brain networks more stable in individuals with higher cognitive abilities (2019, October 15) retrieved 24 April 2024 from <https://medicalxpress.com/news/2019-10-brain-networks-stable->

[individuals-higher.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.