

Researchers investigate why certain brains are more vulnerable to addiction

October 9 2019, by Jenesse Miller



Credit: CC0 Public Domain

In a report published Tuesday, researchers in neuroscience, addiction and behavioral health laid out the latest research into what makes individuals vulnerable to substance abuse and dependency, along with new discoveries that may hold the key for successful prevention and treatment for those addicted to opioids and other drugs.

The authors noted that, although the use of psychoactive drugs has occurred in almost every society in human history, fatalities from the more recent wave of [opioid](#) use are unprecedented. More Americans are dying from overdoses than from car accidents.

Their study, published in *Psychological Science in the Public Interest*, makes the case that understanding the neurobiological mechanisms underlying drug-seeking behaviors is critical, as is finding evidence-based prevention strategies for opioid abuse in particular and [substance abuse](#) in general.

"Addiction is a disease of decision-making," said lead author Antoine Bechara, a professor of psychology at the USC Dornsife College of Letters, Arts and Sciences and an expert in the neuroscience of decision-making, addiction and substance abuse. "The majority of people have intact brain mechanisms of decision-making that keep them resilient to succumbing to an addiction. The question is, who is more vulnerable and how do we best determine that?"

Bechara and his colleagues believe advancements in brain science can help identify those vulnerable individuals before they're exposed to addictive substances, as well as determine how to treat anyone who has become addicted.

The role of the prefrontal cortex in addiction

The authors observed that, during more than three decades of research on drug use, little attention was paid to the importance of the prefrontal [cortex](#)—which is key for self-regulation, long-term goal-setting, impulse control and the ability to predict consequences of behavior—in deciding to take drugs in the first place. As such, they've brought together evidence showing the most vulnerable have abnormal functioning in neurocognitive systems.

In particular, a weak prefrontal cortex—resulting either from [genetic factors](#) such as different levels of neurotransmitters in the brain or from developmental factors like early head injury—may influence [addiction](#). The prefrontal cortex is very susceptible to even mild traumatic head injuries; early child abuse can also impact the development of the prefrontal cortex.

"There are several factors that create the situation where the prefrontal cortex is suboptimal or weak, and the decision-making capacity doesn't develop normally," said Bechara. "These are people who become more susceptible to becoming addicted not just to opioids but other drugs they have access to."

Two extreme positions contribute to opioid addiction and overdose deaths

Bechara argues that, in response to the opioid crisis, two extreme positions have been taken.

"First, the pharmaceutical companies sold the idea that opioid medications will only be used by people in pain and people won't become addicted," he explained. "That's not true, because you have no way of telling who is susceptible to becoming addicted and who is not."

"The overreaction by doctors is another extreme; because of the fear that everyone is going to be addicted to opioids, they are not prescribing them to people in chronic pain who may need them," he continued.

"There are a lot of people who could benefit from controlled administration of those medications, which work very well to treat pain."

Preventing and treating opioid addiction

The authors expressed the need for a clinical test to screen and sort out those who are more likely to misuse and abuse opioid drugs, such as those with prefrontal cortex dysfunction. For those who are already addicted, they presented research into behavioral approaches—including training designed to increase working memory capacity—that can boost the functions of the prefrontal cortex.

The researchers also proposed the use of procedures like [transcranial magnetic stimulation](#), a noninvasive brain-stimulation technique allowing selective neural stimulation or inhibition. The technique, when used to stimulate the executive decision system of the [prefrontal cortex](#), has been shown to reduce craving or consumption of cocaine, cigarettes and alcohol in prior studies.

The authors acknowledged that more research is needed to better understand those therapies and which will be the most effective.

More information: www.psychologicalscience.org/publications/special-section-features/psychological-science-of-addiction.html

Provided by University of Southern California

Citation: Researchers investigate why certain brains are more vulnerable to addiction (2019, October 9) retrieved 24 April 2024 from <https://medicalxpress.com/news/2019-10-brains-vulnerable-addiction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
