

Cannabis use disorder is declining among young adolescents and young adults

October 30 2019



Credit: CC0 Public Domain

The prevalence of cannabis use disorder decreased in 2002 to 2016 among frequent users, according to a new study conducted at Columbia University Mailman School of Public Health. Changes in social attitudes

and the traits of frequent users may explain the decline, according to researchers. This is one of the first studies to examine the general health profile of people using cannabis daily or almost daily and the trends in the prevalence of cannabis use disorder in this population. The findings are online in the journal *Drug and Alcohol Dependence*.

"Contrary to expectations, the frequency of cannabis use disorder among people reporting daily/almost daily use decreased significantly between 2002-2016, said Silvia Martins, MD, Ph.D., associate professor of Epidemiology at Columbia Mailman School. "The findings contradict the predominating hypothesis that the prevalence of DSM-IV CUD would be stable, or increase, among those using with this regularity."

Data from the National Survey on Drug Use and Health for 2002-2016 included 22,651 individuals using cannabis 300+ days in the past year. Cannabis use disorder was defined using DSM-IV criteria for cannabis abuse and/or dependence. Age categories included: 12-17, 18-25, and 26+ and older.

From 2002-2016, the prevalence of cannabis use disorder among people reporting daily or almost daily use decreased across all [age groups](#)—by 27 percent in adolescents, by 30 percent in ages 18-25, and by 37.5 percent for those age 26 and older.

"There could be several reasons behind these declining rates," noted Martins, who is also director, Substance Use Epidemiology Unit at Columbia. "First, the new national cannabis policy environment, with 33 states legalizing medical use and 10 states allowing recreational use of cannabis may have played a role in reducing stigma and perceptions of risk associated with cannabis use. "Secondly, increasing legalization may also be associated with changes in [social attitudes](#) resulting in fewer conflicts with relatives and friends around cannabis use."

For all age groups, there was no evidence of any significant reductions in the perceived need for mental [health](#) treatment among individuals using cannabis regularly (daily/near daily), or the prevalence of health problems as indicated by doctors.

The researchers also did not find evidence of significant reductions in prevalences of past-year health problems when examining health clusters separately including mental health, respiratory, digestive, cardiovascular, and infectious diseases health problems.

In contrast, there were significant decreases in self-reported driving under the influence of illegal drugs including alcohol across all age groups, ranging from a 26 percent, 29 percent and 38 percent change in adolescents, those 18-24 and age 26 and older, respectively.

More information: Julian Santaella-Tenorio et al, Cannabis use disorder among people using cannabis daily/almost daily in the United States, 2002–2016, *Drug and Alcohol Dependence* (2019). [DOI: 10.1016/j.drugalcdep.2019.107621](#)

Provided by Columbia University's Mailman School of Public Health

Citation: Cannabis use disorder is declining among young adolescents and young adults (2019, October 30) retrieved 25 April 2024 from <https://medicalxpress.com/news/2019-10-cannabis-disorder-declining-young-adolescents.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--