

Ebola virus now squeezed into 'corner' of DR Congo: WHO

October 10 2019



Credit: CC0 Public Domain

Efforts to halt an Ebola epidemic in the Democratic Republic of Congo have made "significant progress", with the virus now contained to a far smaller and mainly rural area, the World Health Organization (WHO)

said Thursday.

"We have put the virus in the corner," Michael Ryan, the executive director of WHO's Health Emergencies Programme, told reporters in Geneva.

"I believe we have really squeezed the virus into a much smaller geographical area," he said.

Ebola is now essentially only transmitting within an area of eastern DRC between Mambasa, Komanda, Mandima and Beni, he said.

DRC's latest Ebola epidemic, which began in August 2018, has killed 2,144 people, making it the second deadliest outbreak of the virus, after the West Africa pandemic of 2014-2016.

At the height of the latest outbreak, 207 "health zones" were affected by Ebola, a figure that now stands at only 27, Ryan said.

But he stressed that despite a "much lower level of transmission", the danger was not over.

"The fact that it is a smaller space is positive, but ... the disease has moved into more rural and more insecure areas," he warned.

Ebola fighters have been hindered by militia attacks in eastern DRC, as well as by resistance in communities to some of the methods used to rein in the virus.

'Kill the virus'

"Containing a virus is a different prospect than to eliminate that virus from [human populations](#)," Ryan said.

"We have significantly contained the virus in a much smaller [geographical area](#)," he said. "Now we have to kill the [virus](#)."

Much of the controversy surrounding the response has centered on the use of vaccines.

More than 236,000 people living in active Ebola transmission zones have received an as-yet unlicensed vaccination produced by the pharma giant Merck.

And DRC is preparing to begin introducing a second experimental Ebola [vaccine](#), produced by Johnson & Johnson, to protect those living outside of direct transmission zones.

Ryan said WHO and its partners were talking to communities to explain the differences between the vaccines and say who was entitled to which vaccine and why.

"The two vaccines are complementary. They are designed to do different things in different populations," he said.

The WHO has used a "ring" vaccination approach with the Merck jab, providing it only to the circle of people known to have been in contact with Ebola patients, and their known contacts.

In June, it expanded the access to a "third ring" of people who had possibly had contact with contacts of Ebola patients.

The J&J vaccine, which requires two doses 56 days apart to generate immunity, will meanwhile only be given in a preventative capacity, for instance in [large cities](#) to ensure transmission does not begin there.

It "will provide a sort of firewall", Ryan said.

In July, the WHO declared the Ebola epidemic a "public health emergency of international concern", a rare designation used only for the gravest epidemics.

That move came on the heels of an Ebola case in the city of Goma, one of the country's biggest urban centres.

Containing a lethal contagious disease in cities is invariably far more challenging than in rural settings, given the greater task of tracing and isolating people who are at risk.

The WHO is set to convene its Health Regulations Emergency Committee next week to assess whether the emergency designation should still stand.

© 2019 AFP

Citation: Ebola virus now squeezed into 'corner' of DR Congo: WHO (2019, October 10)
retrieved 19 April 2024 from

<https://medicalxpress.com/news/2019-10-ebola-virus-corner-dr-congo.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--