


Fresh insights could lead to new treatments for liver disease

October 9 2019



Credit: CC0 Public Domain

The fight against liver disease could be helped by the discovery of cells that cause liver scarring.

Scientists have identified new sub-types of cells that, when they interact, accelerate the scarring process in diseased livers.

Experts hope that by understanding more how these cells behave, new treatments can be developed more quickly for liver diseases.

One in five people in the UK are at risk of developing liver disease. It is predicted to become the most common cause of premature death in the UK.

It can occur as a result of a number of conditions such as obesity, alcohol excess, viral infections, autoimmune diseases or genetic disorders.

Long-term damage leads to the formation of scar tissue within the liver, eventually causing liver failure. There are currently no treatments available to prevent or reverse this.

Scientists from the University of Edinburgh used a new technology called single cell RNA sequencing to study liver scarring in [high definition](#).

They discovered sub-types of three key cells: [white blood cells](#) called macrophages, [endothelial cells](#)—which line blood vessels—and scar-forming cells known as myofibroblasts.

The study, published in *Nature*, was funded by the Wellcome Trust, Medical Research Council, Guts UK, Children's Liver Disease Foundation, AbbVie pharmaceuticals, Tenovus Scotland, British Heart Foundation, and NIHR.

This discovery also contributes to the Human Cell Atlas initiative—a global programme designed to create a comprehensive reference map of all cells in the [human body](#).

Dr. Prakash Ramachandran, an MRC clinician scientist at the University of Edinburgh's Centre for Inflammation Research and Consultant Liver

Specialist at the Royal Infirmary of Edinburgh said: "Identifying new treatments for liver scarring is critical to tackling the epidemic of liver disease that we are currently facing. For the first time, we now have an in-depth understanding of how [cells](#) behave and talk to each other in diseased livers and, importantly, how we might block their activity as a treatment for liver scarring."

Professor Neil Henderson, a Wellcome Trust Senior Research Fellow at the Centre for Inflammation Research and Consultant Liver Specialist at the Royal Infirmary of Edinburgh, said: "Using this new technology has allowed us to study human liver scarring in high definition for the first time. We hope that this new cutting-edge approach will rapidly accelerate the discovery of much-needed new treatments for patients with liver [disease](#)."

Dr. Sarah Teichmann, from the Wellcome Sanger Institute, University of Cambridge and co-chair of the Human Cell Atlas Organising Committee, added: "The discovery of three new sub-types of cell involved in human [liver](#) scarring is another important moment in the fight against [liver disease](#) and for the Human Cell Atlas initiative. The data is now publicly available for other researchers to use and will be a great resource as we attempt to tackle this epidemic."

More information: Resolving the fibrotic niche of human liver cirrhosis at single-cell level, *Nature* (2019). [DOI: 10.1038/s41586-019-1631-3](#) , [nature.com/articles/s41586-019-1631-3](#)

Provided by University of Edinburgh

Citation: Fresh insights could lead to new treatments for liver disease (2019, October 9) retrieved 10 April 2024 from

<https://medicalxpress.com/news/2019-10-fresh-insights-treatments-liver-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.