

## Adaptive human immunity depends on the factor responsible for the formation of white blood cells

October 31 2019



Credit: IKBFU



Granulocyte colony-stimulating factor (G-CSF), which causes division and differentiation of bone marrow cells depending on the body's need for leukocytes, has a significant regulatory effect not only on innate, but also on adaptive, immunity. That's what scientists from the IKBFU and Research Institute of Fundamental and Clinical Immunology have found. The scientists have published an article regarding this research in *Growth Factors* journal.

G-CSF plays a key role in the formation of granulocytes in the <u>bone</u> marrow—the most numerous group of white blood <u>cells</u> (leukocytes). This is one of the most important immunological processes, since white blood cells form the basis of the body's anti-infection protection. G-CSF can be produced by cells of the bone marrow, blood vessels, <u>connective</u> <u>tissue</u>, as well as astrocytes that perform a structural function in the nervous system. Moreover, depending on the situation, the <u>human body</u> produces a different amount of G-CSF. This factor acts on stem and myeloid cells of the bone marrow, combining with special receptor proteins on their surface—CD114. This binding enhances <u>cell growth</u> and stimulates the transformation of cells into granulocytes, which have a high non-specific ability to capture and digest viruses and bacteria.

The authors of the study, supported by a grant from the Federal Commission of the Ministry of Education and Science, suggested that G-CSF could be an important tool for regulating the long-term adaptation of immunity to various environmental conditions. To test this, scientists took vein blood from sixteen healthy men and women from 21 to 40 years old. Macrophage cells responsible for the development of adaptive immune responses were isolated from their blood.

The authors then studied the effect that G-CSF exerts on these cells. As a result, it was found that it exerts the most significant stimulating effect on the production of macrophages of IL-6 (interleukin-6), a cytokine involved in the adaptation of various body tissues to the environment. In



addition, G-CSF decreased the number of receptors for proinflammatory cytokine interferon-gamma on macrophages and restrained the development of tissue-damaging inflammation. Based on these data, it can be assumed that G-CSF continuously regulates not only adaptive immunogenesis, but also participates in recovery processes.

Chief Scientist of the IKBFU Center for Medical Biotechnology, Viktor Seledtsov commented on the research results: "Now G-CSF is used mainly for the treatment of leukopenia which is an insufficient amount of <u>white blood cells</u>. However, in the future, it can be used both for the correction of adaptive inflammatory and for the stimulation of regenerative processes in damaged tissues of the body."

**More information:** V. I. Seledtsov et al, Granulocyte colonystimulating factor downregulates interferon-gamma receptor expression and stimulates interleukin-6 production in activated human macrophages, *Growth Factors* (2019). <u>DOI:</u> <u>10.1080/08977194.2019.1662418</u>

## Provided by Immanuel Kant Baltic Federal University

Citation: Adaptive human immunity depends on the factor responsible for the formation of white blood cells (2019, October 31) retrieved 1 May 2024 from <u>https://medicalxpress.com/news/2019-10-human-immunity-factor-responsible-formation.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.