

# Hypoglycaemia prevention could cut hospital stays

October 2 2019

---



Credit: CC0 Public Domain

Preventing in-patients with diabetes from developing hypoglycaemia could dramatically reduce the length of time they spend in hospital—according to new research from the University of East Anglia

and Cambridge University Hospitals NHS Foundation Trust.

Hypoglycaemia happens when blood glucose levels decrease to below [normal levels](#).

The researchers reviewed 15 international studies, comparing outcomes for people with diabetes who were exposed to [hypoglycaemia](#) in [hospital](#) with those with diabetes who were not.

Their findings show that people with diabetes, who were exposed to hypoglycaemia during their admission, had an increased average length of stay of 4.1 days compared to those with diabetes who did not become hypoglycaemic during their admission.

Andrea Lake, a diabetes nurse specialist at Cambridge University Hospitals NHS Foundation Trust (CUH), led the study as part of her master's degree in [clinical research](#) at UEA, funded by the National Institute for Health Research (NIHR).

She said: "It's really important that we understand the impact hypoglycaemia can have on our inpatients' recovery. Our review found not only longer hospital stays for those exposed to hypoglycaemia, but the risk of inpatient mortality almost doubled in this group.

"Although it is impossible to tell from the studies reviewed whether a hypo is the cause of this increase or simply a marker of someone who is at greater risk, understanding the association means we can start to deal with it.

"If we could reduce the chances of it happening in the first place that would be really beneficial for our patients. We need to be considering the risk factors for hypoglycaemia, acting upon them and notifying the appropriate people.

"There are also some simple steps our patients can take to prevent a hypo when they're in hospital. At home or when you're feeling well it's easy to stick to your usual routine. When you're admitted to hospital you might miss your regular snack, or your hypo treatment might have been popped into a fridge, which is kept locked.

"If you're being admitted, bring your normal equipment with you, talk to the team caring for you, and tell them if you're eating less as this means you'll require less insulin. Keep your regular hypo treatment at your bedside along with your glucose meter and if you don't feel great then tell somebody, there may even be an inpatient diabetes team who can assist.

"As part of Hypo Awareness Week we're asking people to become hypo prevention champions. Whether you have diabetes or not, knowing what to look for and what to do can make a real difference. I'm really pleased that the [research paper](#) has shown just how important it is to keep on top of glucose levels when you're in hospital but really it's something you need to be aware of wherever you are."

"The effect of hypoglycaemia during hospital admission on health-related outcomes for people with [diabetes](#): a systematic review and meta-analysis' is published in journal *Diabetic Medicine* to coincide with Hypo Awareness Week, which runs from September 30 until October 6.

**More information:** A. Lake et al. The effect of hypoglycaemia during hospital admission on health-related outcomes for people with diabetes: a systematic review and meta-analysis, *Diabetic Medicine* (2019). [DOI: 10.1111/dme.14115](https://doi.org/10.1111/dme.14115)

Provided by University of East Anglia

Citation: Hypoglycaemia prevention could cut hospital stays (2019, October 2) retrieved 27 April 2024 from <https://medicalxpress.com/news/2019-10-hypoglycaemia-hospital.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.