

## **Researchers use NHANES to search for hypothyroid-sleep apnea link**

October 2 2019

Several past studies have sought to determine if a link exists between thyroid disorders like hypothyroidism, a decrease in thyroid hormone levels, and sleep apnea, a potentially dangerous condition in which a sleeping person's breathing repeatedly stops and starts. However, results from those studies were generally based upon clinical observations that were limited by small population samples and produced often conflicting results.

To more completely understand if such a link may exist, a team from Texas Tech University Health Sciences Center (TTUHSC) embarked upon new research using data mined from the National Health and Nutrition Examination Survey (NHANES). The team included internal medicine resident Subhanudh Thavaraputta, M.D.; Jeff Dennis, Ph.D., an assistant professor for TTUHSC's Department of Public Health who specializes in studying population health; resident Passisd Laoveeravat, M.D., Professor Kenneth Nugent, M.D., and Associate Professor Ana M. Rivas, M.D., from TTUHSC's Department of Internal Medicine. Their study, "Hypothyroidism and its Association with Sleep Apnea Among Adults in the United States: NHANES 2007-2008," was published in July by the *Journal of Clinical Endocrinology & Metabolism*.

The NHANES is a biennial survey conducted by the Centers of Disease Control and Prevention to generally evaluate the health of children and adults in the U.S. In addition to providing a significantly larger sample size, the NHANES provides a cross-sectional sample of the noninstitutionalized U.S. adult population. It includes a detailed



demographic and behavioral questionnaire, a physical examination, laboratory testing and a list of all prescription medications used by the respondent.

Dennis and Thavaraputta used the 2007-2008 NHANES because it is the only one that tested respondents for thyroid stimulating hormone (TSH) levels and included a detailed sleep questionnaire that specifically asked respondents to list any sleep disorder for which they had been diagnosed.

"We had to use those two years to get the exact data points that we wanted," Dennis said. "I always use the most current data when we can, but we were somewhat constrained here, and this is the one NHANES where we could see both TSH levels and sleep disorder information in the same place. We don't have any reason to think that would have changed drastically between the 2007-2008 NHANES and now. What we found should be reasonably consistent over time."

Thavaraputta, the study's lead author, said responses from 5,515 of the 10,000 respondents who participated in the 2007-2008 NHANES were ultimately included in the TTUHSC research. After analyzing those results, he and Dennis estimate that the prevalence for hypothyroidism among U.S. adults is 9.47%, which represents approximately 19.6 million people.

The results also indicate that individuals diagnosed with hypothyroidism are 1.88 times more likely to develop <u>sleep apnea</u>. Hypothyroid patients who were undergoing hormone replacement therapy at the time of the NHANES evaluation were estimated to be 2.51 times more likely to have a sleep apnea diagnosis, regardless of their TSH level.

"We hope that our findings will raise awareness and concern among physicians regarding this association," Thavaraputta said. "In the future, if this relationship is established, assessing the sleep qualities/problems



in hypothyroid patients might be beneficial to improve the patients' standard of care, quality of life and treatment outcomes."

Dennis said the study employed the NHANES to help bridge the public/population health perspective with known clinical insights to more definitively determine if the link between thyroid disorders and sleep apnea is relevant at a clinical level.

In addition, Dennis said the abundant NHANES data provides researchers with opportunities to formulate estimates about population subgroups. In this case, the 2007-2008 NHANES data allowed the TTUHSC team to estimate that up to 2.4% of the U.S. adult population may have undiagnosed hypothyroidism, and that nearly 20% of those diagnosed with hypothyroidism have TSH levels above the normal range.

Working with clinicians like Thavaraputta to build these types of collaborative bridges is a major reason Dennis came to TTUHSC more than four years ago.

"The clinician may have ideas about the patterns they see, but they don't really have time to step back and investigate whether or not this is happening on a national level or with specific groups," he said. "I think that's the huge value of a health sciences center like TTUHSC. It puts people in those different realms together so we can collaborate and hopefully create something that neither of us necessarily would have come up with on our own. I think, ideally, that's the value of what we ended up putting together."

Dennis said the interplay between the clinical and the <u>population health</u> perspectives also was important to the investigation because the research team was unable to clinically diagnose or otherwise verify the sleep disorder information reported by the 5,500-plus individuals whose NHANES responses were used in the study. In the end, he believes that



interplay allowed the TTUHSC team to suggest an association does exist between thyroid disorders and sleep apnea, though they still can't explain why the link occurs.

"We are not suggesting a causal link, but we hope the observed association from our research adds one more piece to the puzzle and sparks further investigation," Dennis said. "I think what we've added could provide value to future clinical and population studies that will continue looking at this potential link."

Provided by Texas Tech University

Citation: Researchers use NHANES to search for hypothyroid-sleep apnea link (2019, October 2) retrieved 4 May 2024 from https://medicalxpress.com/news/2019-10-nhanes-hypothyroid-sleep-apnea-link.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.