


Researchers show path of an environmental estrogen through the womb using new technology

October 10 2019



The food estrogen zearalenone migrates through the placenta, as researchers were able to demonstrate for the first time. Credit: University of Vienna

The human foetus is considered to be particularly sensitive to environmental contaminants. A team led by Benedikt Warth from the Faculty of Chemistry at the University of Vienna and Tina Bürki from

the Swiss Materials Science and Technology Institute, Empa, has now been able to demonstrate for the first time how the widespread food estrogen zearalenone behaves in the womb. Using a new analytical method, it was shown that the xenoestrogen migrates through the placenta and is partially converted to other harmful substances. The study was published in the current issue of the renowned journal *Environmental Health Perspectives*.

Xenoestrogens are absorbed through the environment, especially through food. As estrogen-like substances, they can have a profound effect on the body's hormonal balance. Zearalenone, a widespread food estrogen, is produced by fungi (*Fusarium* species) and enters the body mainly through the diet of bread, cereals and other grain-based products.


"The placental barrier offers the unborn child a certain degree of protection against bacteria, viruses and some foreign substances including some drugs or environmental toxins absorbed by the body. But, as we were able to show for the first time, zearalenone migrates through the placental barrier" says first author Benedikt Warth from the Institute of Food Chemistry and Toxicology at the University of Vienna.

Human placenta models

The path of zearalenone through the womb was demonstrated in experiments with fully functional placentas that were donated after planned Caesarean sections: "The use of human placentas is crucial to achieve predictive results for the transport and metabolism of zearalenone," says Tina Bürki from Empa in St. Gallen. "This is due to considerable species-specific differences in the structure, function and metabolic capacity of the [placenta](#)."

The researchers measured the concentrations of zearalenone in the tissue of the placenta itself as well as in a nutrient solution before entry and

after exit from the placenta, and therefore within the influence of the foetus. At the same time, they were able to study the metabolic products produced by enzymes in the placenta.



Benedikt Warth and master student Karin Preindl, both from the Faculty of Chemistry at the University of Vienna, recently presented a new analytical approach that allows the simultaneous measurement of more than 50 different xenoestrogens in various biological samples. Credit: University of Vienna

Body's metabolism promotes effect

"If we ingest environmental contaminants, they are usually detoxified in the body via our metabolism and then excreted. However, there are also enzymes that make these substances even more bioactive," says Bürki. The researchers showed that the placenta produces metabolic products

that are about a factor 70 more estrogenic than zearalenone. Therefore, even small concentrations could have a greater impact on the embryo than previously assumed.

"This finding should influence future risk assessments, even if the threshold limit values in baby food and breast milk substitutes are already more strictly regulated than for other products and the EU has introduced the world's most protective regulatory limits," explains Benedikt Warth.

The body's hormonal balance is highly sensitive. It is assumed that early life exposure to xenoestrogens may have a significant impact on health outcomes later in life. This could include various diseases such as breast or uterus cancers, but also on other symptoms such as premature puberty or infertility. According to the authors, until further research is available, the best advice is a well-balanced diet in order to avoid high exposures.

Innovative technology as the basis

Only recently, the team around Benedikt Warth, Doris Marko and Karin Preindl from the University of Vienna had provided the pioneering basis for the current work in a study in *Analytical Chemistry*: They presented an analytical method by which more than 50 different xenoestrogens can be measured simultaneously in 20 minutes in various biological samples including urine, blood serum and breast milk. To date, these environmental contaminants were only able to be measured one at a time.

"Our method includes practically all major contaminants that affect the estrogenic system. This also includes various other substances that are currently subject of intense discussion, such as bisphenol A, pesticides or plasticizers," Warth says. With the new method, the researchers hope to be able to better investigate early life co-exposure patterns and

combinatorial effects of [environmental contaminants](#) in the human body in the future.

More information: Benedikt Warth et al. Transfer and Metabolism of the Xenoestrogen Zearalenone in Human Perfused Placenta, *Environmental Health Perspectives* (2019). [DOI: 10.1289/EHP4860](https://doi.org/10.1289/EHP4860)

Karin Preindl et al. A Generic Liquid Chromatography–Tandem Mass Spectrometry Exposome Method for the Determination of Xenoestrogens in Biological Matrices, *Analytical Chemistry* (2019). [DOI: 10.1021/acs.analchem.9b02446](https://doi.org/10.1021/acs.analchem.9b02446)

Provided by University of Vienna

Citation: Researchers show path of an environmental estrogen through the womb using new technology (2019, October 10) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-10-path-environmental-estrogen-womb-technology.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.