

Simple test predicts older heart attack patients' independence, function

October 11 2019, by Brita Belli



Credit: stock.adobe.com

A study just published in *JAMA Internal Medicine* finds that hospitals can administer a simple walking test to evaluate the likelihood of functional decline in older adults following a heart attack. Researchers say the nationwide study could have important implications for how patient care following heart attacks is managed, improving independence

in older adults.

The research was done as part of the SILVER-AMI Study, headquartered at Yale, which focuses exclusively on patients age 75 and over who experienced a [heart attack](#) in order to discover whether collecting information on aging-related conditions improves care and outcomes.

Heart attacks (acute myocardial infarctions, or AMIs) are one of the leading causes of death in the developed world, and more than one-third of heart attacks occur in adults aged 75 and over. The health impacts don't end once patients leave the [hospital](#)—some 30 percent of patients experience losses in function and independence that can last for months.

The study, by Yale epidemiologist Alexandra Hajduk and others, involved 2,587 patients across 94 academic and [community hospitals](#) in the U.S. Patients were given a brief walking assessment called the "Timed Up and Go" or TUG. During the assessment, patients were asked to rise from a seated position, walk a short distance, turn, walk back, and sit back down. Researchers found a strong correlation between poor TUG performance and new limitations in patients' ability to take care of themselves, with difficulty bathing being the most common, followed by difficulty dressing, getting in and out of a chair, and walking around the home.

"This test was able to predict with high accuracy whether patients will lose independence in some of the most important tasks needed to live independently," Hajduk says, noting that these findings could lead to important improvements in the way [patient care](#) is managed following treatment for heart attacks.

"It's increasingly recognized that getting people up and moving in the hospital is important," she says. "But based on these findings, hospital

staff may be able to proactively apply preventive treatment or tailor rehabilitation to a patient's needs, whether that is a physical therapy consultation, [home health care](#), or enhanced [social support](#)."

In the past, says Hajduk, studies of hospitalized older adults have focused largely on two areas: hospital readmission and mortality. But, she says, older adults are most interested in preserving their functional independence—maintaining their ability to perform normal daily tasks as long as they are able. "For many [older adults](#), preserving their independence is the most important goal," Hajduk says.

More information: Alexandra M. Hajduk et al. Association Between Mobility Measured During Hospitalization and Functional Outcomes in Older Adults With Acute Myocardial Infarction in the SILVER-AMI Study, *JAMA Internal Medicine* (2019). [DOI: 10.1001/jamainternmed.2019.4114](#)

Provided by Yale University

Citation: Simple test predicts older heart attack patients' independence, function (2019, October 11) retrieved 6 May 2024 from <https://medicalxpress.com/news/2019-10-simple-older-heart-patients-independence.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--