

# Some 15,000 Spaniards may be unaware they are suffering from the rare bone disease hypophosphatasia

October 31 2019

---



Credit: CC0 Public Domain

Scientists from the research group CIBERFES and the University of Granada (UGR) have implemented a protocol for effectively detecting hypophosphatasia and have discovered two new genetic mutations

associated with this disease, which is often confused with other bone ailments. Extrapolating the data from their study, published in *Scientific Reports*, they estimate that between 4,000 and 15,000 Spaniards may be undiagnosed

In adults, hypophosphatasia causes recurrent fractures and chronic pain. In the [child population](#) this disease can be much more serious and can even cause premature death, neurological disorders, and respiratory problems.

Hypophosphatasia (HPP) is a rare and serious hereditary disease that is potentially life-threatening and is caused by one or more mutations in the gene coding for [alkaline phosphatase](#) (ALPL). It is frequently misdiagnosed.

To progress toward its correct clinical evaluation, researchers from the Centre for Biomedical Research in Fragility and Healthy Ageing (CIBERFES) of the research group led by UGR researcher Dr. Manuel Muñoz Torres in Granada's Biohealth Research Institute of Granada (San Cecilio University Hospital) have implemented a protocol to assess the real proportion of patients affected by this disease. They have also validated the phenotype associated with the two new mutations identified, by means of bioinformatics studies and in vitro assays at the cellular level.

They concluded that almost half of the patients in the study had not been correctly diagnosed and that, extrapolating the data to the Spanish population, there could be as many as 15,000 cases potentially undetected.

Researchers Cristina and Beatriz García Fontana explain that this study, published in *Scientific Reports*, "provides evidence that it is essential to perform a correct clinical assessment to make a correct diagnosis of this

disorder and provide suitable treatment to those patients affected." Due to its low prevalence, this metabolic disorder is usually misdiagnosed and is often confused with more common bone disorders such as osteoporosis, leading to erroneous treatments that could actually worsen the prognosis.

Scientists in the CIBERFES research group estimate that the prevalence in Spain could be twice that quantified in previous studies. To perform this extrapolation, an evaluation of serum levels of total alkaline phosphatase was carried out on a sample of 78,590 patients (76,083 adults and 2,507 children) over the course of 2016. Among the [adult population](#), 1,907 people were identified as having ALP levels below the reference range. After studying the clinical histories of each patient, those subjects who did not meet the criteria established in the proposed protocol were eliminated from the study. Finally, out of the 56 individuals selected, 16 adults (81% of whom were women) agreed to participate. Most of these individuals presented no symptoms at all, or they presented mild symptoms but had never been diagnosed with this condition.

"Using the protocol described in our study, we discovered that seven of these 16 patients were affected by hypophosphatasia, which indicates that almost half of the population we studied was not correctly diagnosed in normal clinical practice," observes Dr. García Fontana of CIBERFES. Extrapolating these data to the Spanish population, there could be 4,000 potential cases of hypophosphatasia currently undiagnosed—and there could be many more.

"Taking into account the proportion of patients who did not participate in the study for different reasons (71%), and according to our results, the current estimate of possible cases of HHP undetected in Spain could be as high as 15,000," continues Dr. García Fontana. This implies that the estimated prevalence of mild HPP in Spain would be twice that

previously published for Europe (1/3,100 vs. 1/6,370).

## History of bone fracture and dental abnormalities

Of the patients analysed at the hospital, none presented a family history of HPP or rickets in childhood, but eight patients (seven of them women) did suffer from prevalent fractures, six of them suffered from symptomatic chondrocalcinosis, and five presented dental anomalies. Regarding the biochemical evidence, 10 adults presented lower alkaline phosphatase and higher levels of the PLP substrate compared to the reference values. Of these ten, seven presented mutations of the ALPL gene, two of which had not previously been described.

Hypophosphatasia in adults involves a loss of mineralization, which gives rise to recurrent fractures, femoral fractures, a history of rickets, and musculoskeletal pain, plus certain dental conditions. Among children it is much more serious, and can cause brain damage, respiratory problems, and even premature death in babies.

**More information:** Cristina García-Fontana et al. Epidemiological, Clinical and Genetic Study of Hypophosphatasia in A Spanish Population: Identification of Two Novel Mutations in The Alpl Gene, *Scientific Reports* (2019). [DOI: 10.1038/s41598-019-46004-2](https://doi.org/10.1038/s41598-019-46004-2)

Provided by University of Granada

Citation: Some 15,000 Spaniards may be unaware they are suffering from the rare bone disease hypophosphatasia (2019, October 31) retrieved 26 April 2024 from <https://medicalxpress.com/news/2019-10-spaniards-unaware-rare-bone-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.