

Stress during pregnancy may affect baby's sex, risk of preterm birth

October 14 2019



Credit: CC0 Public Domain

It's becoming well established that maternal stress during pregnancy can affect fetal and child development as well as birth outcomes, and a new study from researchers at Columbia University Vagelos College of Physicians and Surgeons and NewYork-Presbyterian now identifies the

types of physical and psychological stress that may matter most.

The study was published online in the journal *PNAS*, the *Proceedings of the National Academy of Sciences*.

"The womb is an influential first home, as important as the one a child is raised in, if not more so," says study leader Catherine Monk, Ph.D., professor of medical psychology at Columbia University Vagelos College of Physicians and Surgeons and director of Women's Mental Health in the Department of Obstetrics & Gynecology at NewYork-Presbyterian/Columbia University Irving Medical Center.

Because [stress](#) can manifest in a variety of ways, both as a subjective experience and in physical and lifestyle measurements, Monk and her colleagues examined 27 indicators of psychosocial, physical, and lifestyle stress collected from questionnaires, diaries, and daily physical assessments of 187 otherwise healthy pregnant women, ages 18 to 45.

About 17% (32) of the women were psychologically stressed, with clinically meaningful high levels of depression, anxiety, and perceived stress. Another 16% (30) were physically stressed, with relatively higher daily blood pressure and greater caloric intake compared with other healthy pregnant women. The majority (nearly 67%, or 125) were healthy.

Fewer Baby Boys with Mental Stress?

The study suggested that pregnant women experiencing physical and [psychological stress](#) are less likely to have a boy. On average, around 105 males are born for every 100 female births. But in this study, the sex ratio in the physically and psychologically stressed groups favored girls, with male-to-female ratios of 4:9 and 2:3, respectively.

"Other researchers have seen this pattern after social upheavals, such as the 9/11 terrorist attacks in New York City, after which the relative number of male births decreased," says Monk. "This stress in women is likely of long-standing nature; studies have shown that males are more vulnerable to adverse prenatal environments, suggesting that highly stressed women may be less likely to give [birth](#) to a male due to the loss of prior male pregnancies, often without even knowing they were pregnant."

Other Impacts of Stress

- Physically stressed mothers, with higher blood pressure and caloric intake, were more likely to give birth prematurely than unstressed mothers.
- Among physically stressed mothers, fetuses had reduced heart rate-movement coupling—an indicator of slower central nervous system development—compared with unstressed mothers.
- Psychologically stressed mothers had more birth complications than physically stressed [mothers](#).

Social Support Matters

The researchers also found that what most differentiated the three groups was the amount of social support a mother received from friends and family. For example, the more social support a mother received, the greater the likelihood of her having a male baby.

When social support was statistically equalized across the groups, the stress effects on preterm birth disappeared. "Screening for depression and anxiety are gradually becoming a routine part of prenatal practice," says Monk. "But while our study was small, the results suggest enhancing social support is potentially an effective target for clinical intervention."

An estimated 30% of pregnant [women](#) report psychosocial stress from job strain or related to depression and anxiety, according to the researchers. Such stress has been associated with increased risk of premature birth, which is linked to higher rates of infant mortality and of physical and [mental disorders](#), such as attention-deficit hyperactivity disorder and anxiety, among offspring.

How a mother's mental state might specifically affect a fetus was not examined in the study. "We know from animal studies that exposure to high levels of stress can raise levels of stress hormones like cortisol in the uterus, which in turn can affect the fetus," says Monk. "Stress can also affect the mother's immune system, leading to changes that affect neurological and behavioral development in the fetus. What's clear from our study is that maternal [mental health](#) matters, not only for the mother but also for her future child."

The study is titled, "Maternal prenatal stress phenotypes associate with fetal neurodevelopment and [birth outcomes](#)."

More information: Kate Walsh et al., "Maternal prenatal stress phenotypes associate with fetal neurodevelopment and birth outcomes," *PNAS* (2019). www.pnas.org/cgi/doi/10.1073/pnas.1905890116

Provided by Columbia University Irving Medical Center

Citation: Stress during pregnancy may affect baby's sex, risk of preterm birth (2019, October 14) retrieved 5 May 2024 from <https://medicalxpress.com/news/2019-10-stress-pregnancy-affect-baby-sex.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
