


Combined tests can predict kidney injury risk in critically ill children

November 10 2019



Credit: CC0 Public Domain

Combining 2 tests can improve predictions of severe acute kidney injury in children in intensive care. The findings come from a study that will be presented at ASN Kidney Week 2019 November 5-November 10 at the Walter E. Washington Convention Center in Washington, DC.



Two assessments—the Renal Angina Index (RAI) and measurement of urinary Neutrophil Gelatinase Associated Lipocalin (NGAL)—can be used to determine patients' risk of developing severe acute kidney <u>injury</u>. Kelli Krallman (Cincinnati Children's Hospital Medical Center) and her colleagues looked to determine the potential benefits of combining these assessments.

The team's analysis included 627 pediatric intensive care unit (PICU) admissions. The RAI calculated at 12 hours was found to be a significant predictor of the development of severe acute kidney injury during PICU days 2-4. Adding urinary NGAL assessments for those at risk based off a high RAI score improved the prediction significantly. These RAI+/NGAL+ patients were not only at higher risk for severe acute kidney injury, but also for the need of kidney replacement therapy such as dialysis, longer PICU stay, and longer <u>hospital stay</u>.

"Integration of the RAI and urinary NGAL assessments can be used early in the PICU course to identify patients truly at risk for acute <u>kidney</u> injury and its associated morbidity," said Krallman.

More information: Study: "Predicting Severe AKI, Fluid Overload, and Renal Replacement Therapy with the Renal Angina Index in Critically Ill Children"

Provided by American Society of Nephrology

Citation: Combined tests can predict kidney injury risk in critically ill children (2019, November 10) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2019-11-combined-kidney-injury-critically-ill.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.